

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 December 2003 (24.12.2003)

PCT

(10) International Publication Number
WO 03/106426 A1

(51) International Patent Classification⁷: **C07D 217/04,**
A61K 31/47

[US/US]; 115 Cromwell Lane, West Chester, PA 19380 (US).

(21) International Application Number: **PCT/US03/18778**

(74) Agent: **STEVENS, Lauren, L.**; Beyer Weaver & Thomas, LLP, P.O. BOX 778, Berkeley, CA 94704-0778 (US).

(22) International Filing Date: **12 June 2003 (12.06.2003)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:

✓ 60/389,265 ✓ 14 June 2002 (14.06.2002) US
✓ 60/389,779 ✓ 18 June 2002 (18.06.2002) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(71) Applicants (*for all designated States except US*): **TY-TOKINETICS, INC.** [US/US]; 280 East Grand Avenue, South San Francisco, CA 94080 (US). **SMITHKLINE BEECHAM CORPORATION** [US/US]; One Franklin Plaza, P.O. Box 7929, Philadelphia, PA 19103 (US).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **MCDONALD, Andrew** [US/US]; 1554 Green St. Apartment F, San Francisco, CA 94123 (US). **LI, Pu-Ping** [CA/US]; 638 Pilgrim Drive, Foster City, CA 94404 (US). **BERGNES, Gustave** [US/US]; 74 Kathleen Court, Pacifica, CA 94044 (US). **MORGANS, David, J., Jr.** [US/US]; 281 Vista Grande Avenue, Los Altos, CA 94024 (US). **DHANAK, Dashyant** [GB/US]; 1502 West Woodbank Way, West Chester, PA 19380 (US). **KNIGHT, Steven, David**

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/106426 A1

(54) Title: **COMPOUNDS, COMPOSITIONS, AND METHODS**

(57) Abstract: Compounds useful for treating cellular proliferative diseases and disorders by modulating the activity of KSP are disclosed.

COMPOUNDS, COMPOSITIONS, AND METHODS

CROSS-REFERENCE TO RELATED PATENT APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Patent Application number 60/389,265, filed June 14, 2002 and of U.S. Provisional Patent Application number 60/389,779, filed June 18, 2002; each of which is incorporated herein by reference for all purposes.

FIELD OF THE INVENTION

[0002] This invention relates to compounds which are inhibitors of the mitotic kinesin KSP and are useful in the treatment of cellular proliferative diseases, for example cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders, fungal disorders, and inflammation.

BACKGROUND OF THE INVENTION

[0003] Among the therapeutic agents used to treat cancer are the taxanes and vinca alkaloids, which act on microtubules. Microtubules are the primary structural element of the mitotic spindle. The mitotic spindle is responsible for distribution of replicate copies of the genome to each of the two daughter cells that result from cell division. It is presumed that disruption of the mitotic spindle by these drugs results in inhibition of cancer cell division, and induction of cancer cell death. However, microtubules form other types of cellular structures, including tracks for intracellular transport in nerve processes. Because these agents do not specifically target mitotic spindles, they have side effects that limit their usefulness.

[0004] Improvements in the specificity of agents used to treat cancer is of considerable interest because of the therapeutic benefits which would be realized if the side effects associated with the administration of these agents could be reduced. Traditionally, dramatic improvements in the treatment of cancer are associated with identification of therapeutic agents acting through novel mechanisms. Examples of this include not only the taxanes, but also the camptothecin class of topoisomerase I inhibitors. From both of these perspectives, mitotic kinesins are attractive targets for new anti-cancer agents.

[0005] Mitotic kinesins are enzymes essential for assembly and function of the

mitotic spindle, but are not generally part of other microtubule structures, such as in nerve processes. Mitotic kinesins play essential roles during all phases of mitosis. These enzymes are "molecular motors" that transform energy released by hydrolysis of ATP into mechanical force which drives the directional movement of cellular cargoes along microtubules. The catalytic domain sufficient for this task is a compact structure of approximately 340 amino acids. During mitosis, kinesins organize microtubules into the bipolar structure that is the mitotic spindle. Kinesins mediate movement of chromosomes along spindle microtubules, as well as structural changes in the mitotic spindle associated with specific phases of mitosis. Experimental perturbation of mitotic kinesin function causes malformation or dysfunction of the mitotic spindle, frequently resulting in cell cycle arrest and cell death.

[0006] Among the mitotic kinesins which have been identified is KSP. KSP belongs to an evolutionarily conserved kinesin subfamily of plus end-directed microtubule motors that assemble into bipolar homotetramers consisting of antiparallel homodimers. During mitosis KSP associates with microtubules of the mitotic spindle. Microinjection of antibodies directed against KSP into human cells prevents spindle pole separation during prometaphase, giving rise to monopolar spindles and causing mitotic arrest and induction of programmed cell death. KSP and related kinesins in other, non-human, organisms, bundle antiparallel microtubules and slide them relative to one another, thus forcing the two spindle poles apart. KSP may also mediate in anaphase B spindle elongation and focussing of microtubules at the spindle pole.

[0007] Human KSP (also termed HsEg5) has been described (Blangy, et al., *Cell*, 83:1159-69 (1995); Whitehead, et al., *Arthritis Rheum.*, 39:1635-42 (1996); Galgio et al., *J. Cell Biol.*, 135:339-414 (1996); Blangy, et al., *J Biol. Chem.*, 272:19418-24 (1997); Blangy, et al., *Cell Motil Cytoskeleton*, 40:174-82 (1998); Whitehead and Rattner, *J. Cell Sci.*, 111:2551-61 (1998); Kaiser, et al., *JBC* 274:18925-31 (1999); GenBank accession numbers: X85137, NM004523 and U37426), and a fragment of the KSP gene (TRIP5) has been described (Lee, et al., *Mol Endocrinol.*, 9:243-54 (1995); GenBank accession number L40372). *Xenopus* KSP homologs (Eg5), as well as *Drosophila* KLP61 F/KRP1 30 have been reported.


[0008] Mitotic kinesins, including KSP, are attractive targets for the discovery and development of novel antimitotic chemotherapeutics. Accordingly, it is an object

of the present invention to provide compounds, compositions and methods useful in the inhibition of KSP.

SUMMARY OF THE INVENTION

[0009] In accordance with the objects outlined above, the present invention provides compounds that can be used to treat cellular proliferative diseases. The compounds are KSP inhibitors, particularly human KSP inhibitors. The present invention also provides compositions comprising such compounds, and methods utilizing such compounds or compositions, which can be used to treat cellular proliferative diseases

[0010] In one aspect, the invention relates to methods for treating cellular proliferative diseases, and for treating disorders by inhibiting the activity of KSP. The methods employ compounds represented by Formula I:



Formula I

wherein,

R_1 is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl;

R_2 and $R_{2'}$ are independently chosen from hydrogen, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl; or R_2 and $R_{2'}$ taken together form an optionally substituted 3- to 7-membered ring;

R_{12} is selected from the group consisting of optionally substituted imidazolyl,

optionally substituted imidazoliny, $-\text{NHR}_4$; $-\text{N}(\text{R}_4)(\text{COR}_3)$; $-\text{N}(\text{R}_4)(\text{SO}_2\text{R}_{3a})$; and $-\text{N}(\text{R}_4)(\text{CH}_2\text{R}_{3b})$;

R_3 is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, optionally substituted heteroaralkyl, $\text{R}_{15}\text{O}-$ and $\text{R}_{17}\text{-NH}-$;

R_{3a} is chosen from optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, optionally substituted heteroaralkyl, and $\text{R}_{17}\text{-NH}-$;

R_{3b} is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl;

R_4 is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocycl, and optionally substituted heteroaralkyl;

R_5 , R_6 , R_7 and R_8 are independently chosen from hydrogen, optionally substituted alkyl, optionally substituted alkoxy, halogen, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl, alkylsulfonamido, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, optionally substituted aryl and optionally substituted heteroaryl;

R_{15} is chosen from optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl; and

R_{17} is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl;

(Formula I including single stereoisomers and mixtures of stereoisomers);

a pharmaceutically acceptable salt of a compound of Formula I;

a pharmaceutically acceptable solvate of a compound of Formula I; or

a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I.

[0011] In one aspect, the invention relates to methods for treating cellular proliferative diseases and other disorders that can be treated by inhibiting KSP by the administration of a therapeutically effective amount of a compound of Formula I; a pharmaceutically acceptable salt of a compound of Formula I; a pharmaceutically

acceptable solvate of a compound of Formula I; or a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I. Such diseases and disorders include cancer, hyperplasia, restenosis, cardiac hypertrophy, immune disorders, fungal disorders and inflammation. It will be understood that the method can employ one or more of the foregoing compounds.

[0012] In another aspect, the invention relates to compounds useful in inhibiting KSP kinesin. The compounds are selected from compounds having the structures shown above in Formula I; pharmaceutically acceptable salts of a compound of Formula I; pharmaceutically acceptable solvates of a compound of Formula I; and pharmaceutically acceptable solvates of a pharmaceutically acceptable salt of a compound of Formula I. The invention also relates to pharmaceutical compositions comprising: a therapeutically effective amount of a compound of Formula I; a pharmaceutically acceptable salt of a compound of Formula I; a pharmaceutically acceptable solvate of a compound of Formula I; or a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I; and one or more pharmaceutical excipients. It will be understood that the compositions can employ one or more of the foregoing compounds. In another aspect, the composition further comprises a chemotherapeutic agent other than a compound of the present invention.

[0013] In an additional aspect, the present invention provides methods of screening for compounds that will bind to a KSP kinesin, for example compounds that will displace or compete with the binding of a compound of the invention. The methods comprise combining a labeled compound of the invention, a KSP kinesin, and at least one candidate agent and determining the binding of the candidate agent to the KSP kinesin.

[0014] In a further aspect, the invention provides methods of screening for modulators of KSP kinesin activity. The methods comprise combining a compound of the invention, a KSP kinesin, and at least one candidate agent and determining the effect of the candidate agent on the KSP kinesin activity.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0015] As used in the present specification, the following words and phrases are generally intended to have the meanings as set forth below, except to the extent that the context in which they are used indicates otherwise. The following abbreviations and terms have the indicated meanings throughout:

Ac	=	acetyl
BNB	=	4-bromomethyl-3-nitrobenzoic acid
Boc	=	t-butyloxy carbonyl
Bu	=	butyl
c-	=	cyclo
CBZ	=	carbobenzoxy = benzyloxycarbonyl
DBU	=	diazabicyclo[5.4.0]undec-7-ene
DCM	=	dichloromethane = methylene chloride = CH_2Cl_2
DCE	=	dichloroethane
DEAD	=	diethyl azodicarboxylate
DIC	=	diisopropylcarbodiimide
DIEA	=	N,N-diisopropylethylamine
DMAP	=	4-N,N-dimethylaminopyridine
DMF	=	N,N-dimethylformamide
DMSO	=	dimethyl sulfoxide
DVB	=	1,4-divinylbenzene
EEDQ	=	2-ethoxy-1-ethoxycarbonyl-1,2-dihydroquinoline
Et	=	ethyl
Fmoc	=	9-fluorenylmethoxycarbonyl
GC	=	gas chromatography
HATU	=	O-(7-Azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HMDS	=	hexamethyldisilazane
HOAc	=	acetic acid
HOBt	=	hydroxybenzotriazole
Me	=	methyl
mesyl	=	methanesulfonyl
MTBE	=	methyl t-butyl ether

NMO	=	N-methylmorpholine oxide
PEG	=	polyethylene glycol
Ph	=	phenyl
PhOH	=	phenol
PfP	=	pentafluorophenol
Pht	=	phthalyl
PPTS	=	pyridinium p-toluenesulfonate
Py	=	pyridine
PyBroP	=	bromo-tris-pyrrolidino-phosphonium hexafluorophosphate
rt	=	room temperature
sat'd	=	saturated
s-	=	secondary
t-	=	tertiary
TBDMS	=	t-butyldimethylsilyl
TES	=	triethylsilyl
TFA	=	trifluoroacetic acid
THF	=	tetrahydrofuran
TMOF	=	trimethyl orthoformate
TMS	=	trimethylsilyl
tosyl	=	p-toluenesulfonyl
Trit	=	triphenylmethyl

[0016] **Alkyl** is intended to include linear, branched, or cyclic aliphatic hydrocarbon structures and combinations thereof, which structures may be saturated or unsaturated. **Lower-alkyl** refers to alkyl groups of from 1 to 5 carbon atoms, preferably from 1 to 4 carbon atoms. Examples of lower-alkyl groups include methyl-, ethyl-, propyl-, isopropyl-, butyl-, s- and t-butyl and the like. Preferred alkyl groups are those of C₂₀ or below. More preferred alkyl groups are those of C₁₃ or below. **Cycloalkyl** is a subset of alkyl and includes cyclic aliphatic hydrocarbon groups of from 3 to 13 carbon atoms. Examples of cycloalkyl groups include c-propyl-, c-butyl-, c-pentyl-, norbornyl-, adamantyl and the like. **Cycloalkyl-alkyl-** is another

subset of alkyl and refers to cycloalkyl attached to the parent structure through a non-cyclic alkyl-. Examples of cycloalkyl-alkyl- include cyclohexylmethyl-, cyclopropylmethyl-, cyclohexylpropyl-, and the like. In this application, alkyl includes alkanyl-, alkenyl and alkynyl residues; it is intended to include vinyl-, allyl-, isoprenyl and the like. When an alkyl residue having a specific number of carbons is named, all geometric isomers having that number of carbons are intended to be encompassed; thus, for example, "butyl" is meant to include n-butyl-, sec-butyl-, isobutyl and t-butyl-; "propyl" includes n-propyl-, isopropyl-, and c-propyl-.

[0017] **Alkylene-, alkenylene-, and alkynylene-** are other subsets of alkyl-, including the same residues as alkyl-, but having two points of attachment within a chemical structure. Examples of alkylene include ethylene ($-\text{CH}_2\text{CH}_2-$), propylene ($-\text{CH}_2\text{CH}_2\text{CH}_2-$), dimethylpropylene ($-\text{CH}_2\text{C}(\text{CH}_3)_2\text{CH}_2-$) and cyclohexylpropylene ($-\text{CH}_2\text{CH}_2\text{CH}(\text{C}_6\text{H}_{13})-$). Likewise, examples of alkenylene include ethenylene ($-\text{CH}=\text{CH}-$), propenylene ($-\text{CH}=\text{CH}-\text{CH}_2-$), and cyclohexylpropenylene ($-\text{CH}=\text{CHCH}(\text{C}_6\text{H}_{13})-$). Examples of alkynylene include ethynylene ($-\text{C}\equiv\text{C}-$) and propynylene ($-\text{CH}\equiv\text{CH}-\text{CH}_2-$).

[0018] **Cycloalkenyl** is a subset of alkyl and includes unsaturated cyclic hydrocarbon groups of from 3 to 13 carbon atoms. Examples of cycloalkenyl groups include c-hexenyl-, c-pentenyl and the like.

[0019] **Alkoxy or alkoxyl** refers to an alkyl group, preferably including from 1 to 8 carbon atoms, of a straight, branched, or cyclic configuration, or a combination thereof, attached to the parent structure through an oxygen (i.e., the group alkyl-O-). Examples include methoxy-, ethoxy-, propoxy-, isopropoxy-, cyclopropyloxy-, cyclohexyloxy- and the like. **Lower-alkoxy** refers to alkoxy groups containing one to four carbons.

[0020] **Acyl** refers to groups of from 1 to 8 carbon atoms of a straight, branched, or cyclic configuration or a combination thereof, attached to the parent structure through a carbonyl functionality. Such groups may be saturated or unsaturated, and aliphatic or aromatic. One or more carbons in the acyl residue may be replaced by oxygen, nitrogen (e.g., carboxamido), or sulfur as long as the point of attachment to the parent remains at the carbonyl. Examples include acetyl-, benzoyl-, propionyl-, isobutyryl-, oxalyl-, t-butoxycarbonyl-, benzyloxycarbonyl-, morpholinylcarbonyl, and the like. **Lower-acyl** refers to acyl groups containing one

to four carbons.

[0021] **Amino** refers to the group $-NH_2$. The term "substituted amino" refers to the group $-NHR$ or $-NRR$ where each R is independently selected from the group: optionally substituted alkyl-, optionally substituted alkoxy, optionally substituted aminocarbonyl-, optionally substituted aryl-, optionally substituted heteroaryl-, optionally substituted heterocyclyl-, acyl-, alkoxycarbonyl-, sulfanyl-, sulfinyl and sulfonyl-, e.g., diethylamino, methylsulfonylamino, furanyl-oxy-sulfonamino.

[0022] **Aminocarbonyl-** refers to the group $-NR^cCOR^b$, $-NR^cCO_2R^a$, or $-NR^cCONR^bR^c$, where

R^a is optionally substituted C_1 - C_6 alkyl-, aryl-, heteroaryl-, aryl- C_1 - C_4 alkyl-, or heteroaryl- C_1 - C_4 alkyl- group;

R^b is H or optionally substituted C_1 - C_6 alkyl-, aryl-, heteroaryl-, aryl- C_1 - C_4 alkyl-, or heteroaryl- C_1 - C_4 alkyl- group; and

R^c is hydrogen, alkyl-, aryl- or heteroaryl-; and

where each optionally substituted R^b group is independently unsubstituted or substituted with one or more substituents independently selected from C_1 - C_4 alkyl-, aryl-, heterocyclyl-, aryl- C_1 - C_4 alkyl-, heteroaryl- C_1 - C_4 alkyl-, C_1 - C_4 haloalkyl-, $-OC_1$ - C_4 alkyl, $-OC_1$ - C_4 alkylphenyl, $-C_1$ - C_4 alkyl-OH, $-OC_1$ - C_4 haloalkyl, halogen, $-OH$, $-NH_2$, $-NR^cC(NR^b)(NR^bR^c)$ (i.e., guanidine), $-NR^cCR^bNR^bR^c$, $-CNR^cNR^bR^c$, $-C_1$ - C_4 alkyl- NH_2 , $-N(C_1$ - C_4 alkyl)(C_1 - C_4 alkyl), $-NH(C_1$ - C_4 alkyl), $-N(C_1$ - C_4 alkyl)(C_1 - C_4 alkylphenyl), $-NH(C_1$ - C_4 alkylphenyl), cyano, nitro, oxo (as a substituent for heteroaryl), $-CO_2H$, $-C(O)OC_1$ - C_4 alkyl-, $-CON(C_1$ - C_4 alkyl)(C_1 - C_4 alkyl), $-CONH(C_1$ - C_4 alkyl), $-CONH_2$, $-NHC(O)(C_1$ - C_4 alkyl), $-NHC(O)(phenyl)$, $-N(C_1$ - C_4 alkyl) $C(O)(C_1$ - C_4 alkyl), $-N(C_1$ - C_4 alkyl) $C(O)(phenyl)$, $-C(O)C_1$ - C_4 alkyl-, $-C(O)C_1$ - C_4 phenyl-, $-C(O)C_1$ - C_4 haloalkyl-, $-OC(O)C_1$ - C_4 alkyl-, $-SO_2(C_1$ - C_4 alkyl), $-SO_2(phenyl)$, $-SO_2(C_1$ - C_4 haloalkyl), $-SO_2NH_2$, $-SO_2NH(C_1$ - C_4 alkyl), $-SO_2NH(phenyl)$, $-NHSO_2(C_1$ - C_4 alkyl), $-NHSO_2(phenyl)$, and $-NHSO_2(C_1$ - C_4 haloalkyl).

[0023] **Antimitotic** refers to a drug for inhibiting or preventing mitosis, for example, by causing metaphase arrest. Some antitumour drugs block proliferation and are considered antimitotics.

[0024] **Aryl** and **heteroaryl** mean a 5- or 6-membered aromatic or heteroaromatic ring containing 0 or 1-4 heteroatoms, respectively, selected from O, N,

or S; a bicyclic 9- or 10-membered aromatic or heteroaromatic ring system containing 0 or 1-4 (or more) heteroatoms, respectively, selected from O, N, or S; or a tricyclic 12- to 14-membered aromatic or heteroaromatic ring system containing 0 or 1-4 (or more) heteroatoms, respectively, selected from O, N, or S. The aromatic 6- to 14-membered carbocyclic rings include, e.g., phenyl-, naphthyl-, indanyl-, tetralinyl-, and fluorenyl and the 5- to 10-membered aromatic heterocyclic rings include, e.g., imidazolyl-, pyridinyl-, indolyl-, thienyl-, benzopyranonyl-, thiazolyl-, furanyl-, benzimidazolyl-, quinolinyl-, isoquinolinyl-, quinoxalinyl-, pyrimidinyl-, pyrazinyl-, tetrazolyl and pyrazolyl-.

[0025] **Aralkyl-** refers to a residue in which an aryl moiety is attached to the parent structure via an alkyl residue. Examples include benzyl-, phenethyl-, phenylvinyl-, phenylallyl and the like. **Heteroaralkyl-** refers to a residue in which a heteroaryl moiety is attached to the parent structure via an alkyl residue. Examples include furanylmethyl-, pyridinylmethyl-, pyrimidinylethyl and the like.

[0026] **Aralkoxy-** refers to the group -O-aralkyl. Similarly, **heteroaralkoxy-** refers to the group -O-heteroaralkyl-; **aryloxy-** refers to the group -O-aryl-; **acyloxy-** refers to the group -O-acyl-; **heteroaryloxy-** refers to the group -O-heteroaryl-; and **heterocyclyloxy-** refers to the group -O-heterocyclyl (i.e., aralkyl-, heteroaralkyl-, aryl-, acyl-, heterocyclyl-, or heteroaryl is attached to the parent structure through an oxygen).

[0027] **Carboxyalkyl-** refers to the group -alkyl-COOH.

[0028] **Carboxamido** refers to the group -CONR^bR^c, where

R^b is H or optionally substituted C₁-C₆ alkyl-, aryl-, heteroaryl-, aryl-C₁-C₄ alkyl-, or heteroaryl-C₁-C₄ alkyl- group; and

R^c is hydrogen, alkyl-, aryl- or heteroaryl-; and

where each optionally substituted R^b group is independently unsubstituted or substituted with one or more substituents independently selected from C₁-C₄ alkyl-, aryl-, heterocyclyl-, aryl-C₁-C₄ alkyl-, heteroaryl-C₁-C₄ alkyl-, C₁-C₄ haloalkyl-, -OC₁-C₄ alkyl-, -OC₁-C₄ alkylphenyl-, -C₁-C₄ alkyl-OH, -OC₁-C₄ haloalkyl-, halogen, -OH, -NH₂, -C₁-C₄ alkyl-NH₂, -N(C₁-C₄ alkyl)(C₁-C₄ alkyl), -NH(C₁-C₄ alkyl), -N(C₁-C₄ alkyl)(C₁-C₄ alkylphenyl), -NH(C₁-C₄ alkylphenyl), cyano, nitro, oxo (as a substituent for heteroaryl), -CO₂H, -C(O)OC₁-C₄ alkyl-, -CON(C₁-C₄ alkyl)(C₁-C₄ alkyl), -CONH(C₁-C₄ alkyl), -CONH₂,

-NHC(O)(C₁-C₄ alkyl), -NHC(O)(phenyl), -N(C₁-C₄ alkyl)C(O)(C₁-C₄ alkyl),
 -N(C₁-C₄ alkyl)C(O)(phenyl), -C(O)C₁-C₄ alkyl-, -C(O)C₁-C₄ phenyl-,
 -C(O)C₁-C₄ haloalkyl-, -OC(O)C₁-C₄ alkyl-, -SO₂(C₁-C₄ alkyl), -SO₂(phenyl), -
 SO₂(C₁-C₄ haloalkyl), -SO₂NH₂, -SO₂NH(C₁-C₄ alkyl), -SO₂NH(phenyl), -
 NHSO₂(C₁-C₄ alkyl), -NHSO₂(phenyl), and -NHSO₂(C₁-C₄ haloalkyl). Carboxamido
 is meant to include carbamoyl-, lower-alkyl carbamoyl-, benzylcarbamoyl-,
 phenylcarbamoyl-, methoxymethyl-carbamoyl-, and the like.

[0029] **Halogen** or **halo** refers to fluorine, chlorine, bromine or iodine. Fluorine, chlorine and bromine are preferred. Dihaloaryl-, dihaloalkyl-, trihaloaryl etc. refer to aryl and alkyl substituted with the designated plurality of halogens (here, 2, 2 and 3, respectively), but not necessarily a plurality of the same halogen; thus 4-chloro-3-fluorophenyl is within the scope of dihaloaryl-.

[0030] **Heterocyclyl** means a cycloalkyl or aryl residue in which one to four of the carbons is replaced by a heteroatom such as oxygen, nitrogen or sulfur. Examples of heterocycles that fall within the scope of the invention include azetidiny-, imidazolinyl-, pyrrolidinyl-, pyrazolyl-, pyrrolyl-, indolyl-, quinolinyl-, isoquinolinyl-, tetrahydroisoquinolinyl-, benzofuranyl-, benzodioxanyl-, benzodioxyl (commonly referred to as methylenedioxyphenyl-, when occurring as a substituent), tetrazolyl-, morpholinyl-, thiazolyl-, pyridinyl-, pyridazinyl-, piperidinyl-, pyrimidinyl-, thienyl-, furanyl-, oxazolyl-, oxazolinyl-, isoxazolyl-, dioxanyl-, tetrahydrofuranlyl and the like. "N-heterocyclyl" refers to a nitrogen-containing heterocycle. The term heterocyclyl encompasses heteroaryl-, which is a subset of heterocyclyl-. Examples of N-heterocyclyl residues include azetidiny-, 4-morpholinyl-, 4-thiomorpholinyl-, 1-piperidinyl-, 1-pyrrolidinyl-, 3-thiazolidinyl-, piperazinyl and 4-(3,4-dihydrobenzoxazinyl). Examples of substituted heterocyclyl include 4-methyl-1-piperazinyl and 4-benzyl-1-piperidinyl-.

[0031] **A leaving group or atom** is any group or atom that will, under the reaction conditions, cleave from the starting material, thus promoting reaction at a specified site. Suitable examples of such groups unless otherwise specified are halogen atoms, mesyloxy, p-nitrobenzensulphonyloxy and tosyloxy groups.

[0032] **Optional** or **optionally** means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstances occurs and instances in which it does not. For example,

“optionally substituted alkyl” includes “alkyl” and “substituted alkyl” as defined herein. It will be understood by those skilled in the art with respect to any group containing one or more substituents that such groups are not intended to introduce any substitution or substitution patterns that are sterically impractical and/or synthetically non-feasible and/or inherently unstable.

[0033] **Substituted alkoxy** refers to alkoxy wherein the alkyl constituent is substituted (i.e., -O-(substituted alkyl)). One suitable substituted alkoxy group is “polyalkoxy” or -O-(optionally substituted alkylene)-(optionally substituted alkoxy), and includes groups such as -OCH₂CH₂OCH₃, and residues of glycol ethers such as polyethyleneglycol, and -O(CH₂CH₂O)_xCH₃, where x is an integer of about 2-20, preferably about 2-10, and more preferably about 2-5. Another suitable substituted alkoxy group is hydroxyalkoxy or -OCH₂(CH₂)_yOH, where y is an integer of about 1-10, preferably about 1-4.

[0034] **Substituted-** alkyl-, aryl-, and heteroaryl- refer respectively to alkyl-, aryl-, and heteroaryl wherein one or more (up to about 5, preferably up to about 3) hydrogen atoms are replaced by a substituent independently selected from the group: -R^a, -OR^b, -O(C₁-C₂ alkyl)O- (as an aryl substituent), -SR^b, -NR^bR^c, -NR^cC(NR^b)(NR^bR^c) (i.e., guanidine), -NR^cCR^bNR^bR^c, -CNR^cNR^bR^c, halogen, cyano, nitro, -COR^b, -CO₂R^b, -CONR^bR^c, -OCOR^b, -OCO₂R^a, -OCONR^bR^c, -NR^cCOR^b, -NR^cCO₂R^a, -NR^cCONR^bR^c, -CO₂R^b, -CONR^bR^c, -NR^cCOR^b, -SOR^a, -SO₂R^a, -SO₂NR^bR^c, and -NR^cSO₂R^a,

where R^a is an optionally substituted C₁-C₆ alkyl-, aryl-, heteroaryl-, aryl-C₁-C₄ alkyl-, or heteroaryl-C₁-C₄ alkyl- group,

R^b is H or optionally substituted C₁-C₆ alkyl-, aryl-, heteroaryl-, aryl-C₁-C₄ alkyl-, or heteroaryl-C₁-C₄ alkyl- group;

R^c is hydrogen, alkyl-, aryl- or heteroaryl-; and

where each optionally substituted R^a group and R^b group is independently unsubstituted or substituted with one or more substituents independently selected from C₁-C₄ alkyl-, aryl-, heterocycl-yl-, aryl-C₁-C₄ alkyl-, heteroaryl-C₁-C₄ alkyl-, C₁-C₄ haloalkyl-, -OC₁-C₄ alkyl-, -OC₁-C₄ alkylphenyl-, -C₁-C₄ alkyl-OH, -OC₁-C₄ haloalkyl-, halogen, -OH, -NH₂, -C₁-C₄ alkyl-NH₂, -N(C₁-C₄ alkyl)(C₁-C₄ alkyl), -NH(C₁-C₄ alkyl), -N(C₁-C₄ alkyl)(C₁-C₄ alkylphenyl), -NH(C₁-C₄ alkylphenyl), cyano, nitro, oxo (as a substituent for heteroaryl), -CO₂H,

-C(O)OC₁-C₄ alkyl-, -CON(C₁-C₄ alkyl)(C₁-C₄ alkyl), -CONH(C₁-C₄ alkyl), -CONH₂, -NHC(O)(C₁-C₄ alkyl), -NHC(O)(phenyl), -N(C₁-C₄ alkyl)C(O)(C₁-C₄ alkyl), -N(C₁-C₄ alkyl)C(O)(phenyl), -C(O)C₁-C₄ alkyl-, -C(O)C₁-C₄ phenyl-, -C(O)C₁-C₄ haloalkyl-, -OC(O)C₁-C₄ alkyl-, -SO₂(C₁-C₄ alkyl), -SO₂(phenyl), -SO₂(C₁-C₄ haloalkyl), -SO₂NH₂, -SO₂NH(C₁-C₄ alkyl), -SO₂NH(phenyl), -NHSO₂(C₁-C₄ alkyl), -NHSO₂(phenyl), and -NHSO₂(C₁-C₄ haloalkyl).

[0035] **Sulfanyl** refers to the groups: -S-(optionally substituted alkyl), -S-(optionally substituted aryl), -S-(optionally substituted heteroaryl), and -S-(optionally substituted heterocyclyl).

[0036] **Sulfinyl** refers to the groups: -S(O)-H, -S(O)-(optionally substituted alkyl), -S(O)-optionally substituted aryl, -S(O)-(optionally substituted heteroaryl), -S(O)-(optionally substituted heterocyclyl); and -S(O)-(optionally substituted amino).

[0037] **Sulfonyl** refers to the groups: -S(O₂)-H, -S(O₂)-(optionally substituted alkyl), -S(O₂)-optionally substituted aryl, -S(O₂)-(optionally substituted heteroaryl), -S(O₂)-(optionally substituted heterocyclyl), -S(O₂)-(optionally substituted alkoxy), -S(O₂)-optionally substituted aryloxy, -S(O₂)-(optionally substituted heteroaryloxy), -S(O₂)-(optionally substituted heterocyclyloxy); and -S(O₂)-(optionally substituted amino).

[0038] **Pharmaceutically acceptable salts** refers to those salts that retain the biological effectiveness of the free compound and that are not biologically undesirable or unsuitable for pharmaceutical use, formed with a suitable acid or base, and includes pharmaceutically acceptable acid addition salts and base addition salts.

Pharmaceutically acceptable acid addition salts include those derived from inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and those derived from organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

[0039] **Pharmaceutically acceptable base addition salts** include those derived from inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Particular embodiments are the ammonium, potassium, sodium, calcium, and

magnesium salts. Base addition salts also include those derived from pharmaceutically acceptable organic non-toxic bases, including salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, and ethanolamine.

[0040] **Protecting group** has the meaning conventionally associated with it in organic synthesis, i.e. a group that selectively blocks one or more reactive sites in a multifunctional compound such that a chemical reaction can be carried out selectively on another unprotected reactive site and such that the group can readily be removed after the selective reaction is complete. A variety of protecting groups are disclosed, for example, in T.H. Greene and P. G. M. Wuts, *Protective Groups in Organic Synthesis*, Third Edition, John Wiley & Sons, New York (1999), which is incorporated herein by reference in its entirety. For example, a hydroxy protected form is where at least one of the hydroxyl groups present in a compound is protected with a hydroxy protecting group. Likewise, amines and other reactive groups may similarly be protected.

[0041] **Solvate** refers to the compound formed by the interaction of a solvent and a compound of Formula I or salt thereof, and as will be understood by those skilled in the art, is the compound or salt with which a solvent is incorporated, for example, into the crystal structure. Suitable solvates of the compounds of the Formula I or a salt thereof are those formed with a pharmaceutically acceptable solvent, including hydrates (i.e., wherein the solvent is water).

[0042] Many of the compounds described herein contain one or more asymmetric centers (e.g. the carbon to which R_2 and $R_{2'}$ are attached where R_2 differs from $R_{2'}$) and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (R)- or (S)-. The present invention is meant to include all such possible isomers, including racemic mixtures, optically pure forms and intermediate mixtures. Optically active (R)- and (S)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. When the compounds described herein contain olefinic double bonds or other centers of geometric asymmetry, and unless specified otherwise, it is intended that the compounds include both E and Z

geometric isomers. Likewise, all tautomeric forms and rotational isomers are also intended to be included.

[0043] When desired, the R- and S-isomers may be resolved by methods known to those skilled in the art, for example by formation of diastereoisomeric salts or complexes which may be separated, for example, by crystallization; via formation of diastereoisomeric derivatives which may be separated, for example, by crystallization, gas-liquid or liquid chromatography; selective reaction of one enantiomer with an enantiomer-specific reagent, for example enzymatic oxidation or reduction, followed by separation of the modified and unmodified enantiomers; or gas-liquid or liquid chromatography in a chiral environment, for example on a chiral support, such as silica with a bound chiral ligand or in the presence of a chiral solvent. It will be appreciated that where the desired enantiomer is converted into another chemical entity by one of the separation procedures described above, a further step may be required to liberate the desired enantiomeric form. Alternatively, specific enantiomer may be synthesized by asymmetric synthesis using optically active reagents, substrates, catalysts or solvents, or by converting one enantiomer to the other by asymmetric transformation.

Compounds of the Present Invention


[0044] The present invention is directed to a class of novel compounds, that can be described as isoquinolone derivatives, that are inhibitors of one or more mitotic kinesins. By inhibiting mitotic kinesins, but not other kinesins (e.g., transport kinesins), specific inhibition of cellular proliferation is accomplished. While not intending to be bound by any theory, the present invention capitalizes on the finding that perturbation of mitotic kinesin function causes malformation or dysfunction of mitotic spindles, frequently resulting in cell cycle arrest and cell death. According to one embodiment of the invention, the compounds described herein inhibit the mitotic kinesin, KSP, particularly human KSP. In another embodiment, the compounds inhibit the mitotic kinesin, KSP, as well as modulating one or more of the human mitotic kinesins selected from the group consisting of HSET (see, U.S. Patent No. 6,361,993, which is incorporated herein by reference); MCAK (see, U.S. Patent No. 6,331,424, which is incorporated herein by reference); CENP-E (see, PCT Publication No. WO 99/13061, which is incorporated herein by reference); Kif4 (see,

U.S. Patent No. 6,440,684, which is incorporated herein by reference); MKLP1 (see, U.S. Patent No. 6,448,025, which is incorporated herein by reference); Kif15 (see, U.S. Patent No. 6,355,466, which is incorporated herein by reference); Kid (see, U.S. Patent No. 6,387,644, which is incorporated herein by reference); Mpp1, CMKrp, KinI-3 (see, U.S. Patent No. 6,461,855, which is incorporated herein by reference); Kip3a (see, PCT Publication No. WO 01/96593, which is incorporated herein by reference); Kip3d (see, U.S. Patent No. 6,492,151, which is incorporated herein by reference); and RabK6.

[0045] The methods of inhibiting a mitotic kinesin comprise contacting an inhibitor of the invention with a kinesin, particularly a human kinesin, more particularly, human KSP or fragments and variants thereof. The inhibition can be of the ATP hydrolysis activity of the KSP kinesin and/or the mitotic spindle formation activity, such that the mitotic spindles are disrupted. Meiotic spindles may also be disrupted.

[0046] The present invention provides inhibitors of mitotic kinesins, in particular KSP and especially human KSP, for the treatment of disorders associated with cell proliferation. The compounds, compositions and methods described herein can differ in their selectivity and are used to treat diseases of cellular proliferation, including, but not limited to cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders, fungal disorders and inflammation.

[0047] Accordingly, the present invention relates to methods employing compounds represented by Formula I:



Formula I

wherein,

R_1 is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl;

R_2 and $R_{2'}$ are independently chosen from hydrogen, optionally substituted alkyl, optionally substituted alkoxy, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl; or R_2 and $R_{2'}$ taken together form an optionally substituted 3- to 7-membered ring;

R_{12} is selected from the group consisting of optionally substituted imidazolyl, optionally substituted imidazolyl, $-NHR_4$; $-N(R_4)(COR_3)$; $-N(R_4)(SO_2R_{3a})$; and $-N(R_4)(CH_2R_{3b})$;

R_3 is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, optionally substituted heteroaralkyl, $R_{15}O-$ and $R_{17}-NH-$;

R_{3a} is chosen from optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, optionally substituted heteroaralkyl, and $R_{17}-NH-$;

R_{3b} is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl;

R_4 is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heterocyclyl, and optionally substituted heteroaralkyl;

R_5 , R_6 , R_7 and R_8 are independently chosen from hydrogen, optionally substituted alkyl, optionally substituted alkoxy, halogen, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl, alkylsulfonamido, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, optionally substituted aryl and optionally substituted heteroaryl;

R_{15} is chosen from optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl; and

R_{17} is chosen from hydrogen, optionally substituted alkyl, optionally

substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl;

including single stereoisomers and mixtures of stereoisomers;


a pharmaceutically acceptable salt of a compound of Formula I;

a pharmaceutically acceptable solvate of a compound of Formula I;

or a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I. In a particular embodiment, the stereogenic center to which R₂ and R₂' are attached is of the R configuration.


Nomenclature

[0048] The compounds of Formula I can be named and numbered in the manner described below (e.g., using AutoNom version 2.1 in ISIS-DRAW or ChemDraw). For example, the compound:



i.e., the compound according to Formula I where R₁ is benzyl-, R₂ is propyl (or i-propyl), R₂' is hydrogen; R₁₂ is substituted imidazolinyI wherein R₁₀ and R₁₀' are methyl, R₁₁ and R₁₁' are hydrogen and R₁₄ is p-tolyl; R₅, R₆, and R₈ are hydrogen; and R₇ is chloro can be named 2-Benzyl-6-chloro-3-[1-(4,4-dimethyl-2-p-tolyl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-2H-isoquinolin-1-one.

[0049] Likewise, the compound:



i.e., the compound according to Formula I where R_1 is benzyl-, R_2 is propyl (or i - propyl), R_2' is hydrogen; R_{12} is -N(R_4)(COR₃) wherein R_4 is 3-aminopropyl- and R_3 is p-tolyl; R_5 , R_6 , and R_8 are hydrogen; and R_7 is chloro can be named N-(3-Amino-propyl)-N-[1-(2-benzyl-6-chloro-1-oxo-1,2-dihydro-isoquinolin-3-yl)-2-methyl-propyl]-4-methyl-benzamide.

Synthetic Reaction Parameters

[0050] The compounds of Formula I can be prepared by following the procedures described with reference to the Reaction Schemes below.

[0051] Unless specified otherwise, the terms "solvent", "inert organic solvent" or "inert solvent" mean a solvent inert under the conditions of the reaction being described in conjunction therewith [including, for example, benzene, toluene, acetonitrile, tetrahydrofuran ("THF"), dimethylformamide ("DMF"), chloroform, methylene chloride (or dichloromethane), diethyl ether, methanol, pyridine and the like]. Unless specified to the contrary, the solvents used in the reactions of the present invention are inert organic solvents.

[0052] The term "q.s." means adding a quantity sufficient to achieve a stated function, e.g., to bring a solution to the desired volume (i.e., 100%).

[0053] In general, esters of carboxylic acids may be prepared by conventional esterification procedures, for example alkyl esters may be prepared by treating the required carboxylic acid with the appropriate alkanol, generally under acidic conditions. Likewise, amides may be prepared using conventional amidation procedures, for example amides may be prepared by treating an activated carboxylic acid with the appropriate amine. Alternatively, a lower-alkyl ester such as a methyl ester of the acid may be treated with an amine to provide the required amide, optionally in presence of trimethylaluminum following the procedure described in Tetrahedron Lett. 48, 4171-4173, (1977). Carboxyl groups may be protected as alkyl esters, for example methyl esters, which esters may be prepared and removed using conventional procedures, one convenient method for converting carbomethoxy to carboxyl is to use aqueous lithium hydroxide.

[0054] The salts and solvates of the compounds mentioned herein may as required be produced by methods conventional in the art. For example, if an inventive compound is an acid, a desired base addition salt can be prepared by treatment of the free acid with an inorganic or organic base, such as an amine (primary, secondary, or tertiary); an alkali metal or alkaline earth metal hydroxide; or the like. Illustrative examples of suitable salts include organic salts derived from amino acids such as glycine and arginine; ammonia; primary, secondary, and tertiary amines; such as ethylenediamine, and cyclic amines, such as cyclohexylamine, piperidine, morpholine, and piperazine; as well as inorganic salts derived from sodium, calcium, potassium, magnesium, manganese, iron, copper, zinc, aluminum, and lithium.

[0055] If a compound is a base, a desired acid addition salt may be prepared by any suitable method known in the art, including treatment of the free base with an inorganic acid, such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like, or with an organic acid, such as acetic acid, maleic acid, succinic acid, mandelic acid, fumaric acid, malonic acid, pyruvic acid, oxalic acid, glycolic acid, salicylic acid, pyranosidyl acid, such as glucuronic acid or galacturonic acid, alpha-hydroxy acid, such as citric acid or tartaric acid, amino acid, such as aspartic acid or glutamic acid, aromatic acid, such as benzoic acid or cinnamic acid, sulfonic acid, such as p-toluenesulfonic acid, methanesulfonic acid, ethanesulfonic

acid, or the like.

[0056] Isolation and purification of the compounds and intermediates described herein can be effected, if desired, by any suitable separation or purification procedure such as, for example, filtration, extraction, crystallization, column chromatography, thin-layer chromatography or thick-layer chromatography, or a combination of these procedures. Specific illustrations of suitable separation and isolation procedures can be had by reference to the examples hereinbelow. However, other equivalent separation or isolation procedures can, of course, also be used.

Synthesis of the Compounds of Formula I

[0057] The compounds of Formula I can be prepared by following the procedures described in U.S. Patent Application Serial No. 10/366,828 and corresponding PCT Application No. US03/04713; and PCT Publication Nos. WO 01/30768 and WO 01/98278, each of which is incorporated herein by reference for all purposes, and with reference to the Reaction Schemes below.

Brief Description Of Reaction Schemes

[0058] Reaction Scheme 1 illustrates a synthesis of an intermediate (i.e., compounds of Formula 113) in the synthesis of compounds of Formula I.

[0059] Reaction Scheme 2 illustrates a synthesis of compounds of Formula I wherein R_{12} is $-N(R_4)(COR_3)$ from compounds of Formula 113.

[0060] Reaction Scheme 3 shows a synthesis of compounds of Formula I wherein R_{12} is $-N(R_4)(SO_2R_{3a})$ from compounds of Formula 203.

[0061] Reaction Scheme 4 shows a synthesis of compounds of Formula I wherein R_{12} is $-N(R_4)(CH_2R_{3b})$ from compounds of Formula 203.

[0062] Reaction Scheme 5 shows a synthesis of compounds of Formula I wherein R_{12} is optionally substituted imidazolyl from compounds of Formula 113.

[0063] Reaction Scheme 6 shows a synthesis of compounds of Formula I wherein R_{12} is optionally substituted imidazolyl from compounds of Formula 113.

[0064] Reaction Scheme 7 shows a synthesis of compounds of Formula I

wherein R_{12} is optionally substituted imidazoliny from compounds of Formula 113.

[0065] Reaction Scheme 8 shows a synthesis of compounds of Formula I wherein R_{12} is optionally substituted imidazoliny from compounds of Formula 705.

[0066] Reaction Scheme 9 shows a synthesis of compounds of Formula I wherein R_{12} is $-N(R_4)(COR_3)$ and R_3 is $-OR_{15}$, from compounds of Formula 203.

[0067] Reaction Scheme 10 shows a synthesis of compounds of Formula I wherein R_{12} is $-N(R_4)(COR_3)$ and R_3 is $-NHR_{17}$, from compounds of Formula 203.


[0068] Reaction Scheme 11 shows a synthesis of compounds of Formula I wherein R_{12} is optionally substituted imidazolyl.


[0069] Reaction Scheme 12 shows a synthesis of compounds of Formula I wherein R_{12} is optionally substituted imidazoliny from compounds of Formula 113.

Starting Materials

[0070] The optionally substituted benzoic acids of Formula 101 and the other reactants are commercially available, e.g., from Aldrich Chemical Company, Milwaukee, WI or may be readily prepared by those skilled in the art using commonly employed synthetic methodology.

Reaction Scheme 1





Preparation of Compounds of Formula 103

[0071] Referring to Reaction Scheme 1, Step 1, an optionally substituted carboxylic acid of Formula 101 is protected with a suitable protecting group. For example, the following procedure is used to prepare esters of the carboxylic acid. A solution of a compound of Formula 101 and thionyl chloride in a polar aprotic solvent (such as DMF) is warmed gently until the mixture becomes homogeneous. The solution is concentrated. A lower alkanol, such as methanol, is then added. The corresponding, optionally substituted compound of Formula 103 is isolated and purified.

Preparation of Compounds of Formula 105

[0072] Referring to Reaction Scheme 1, Step 2, a compound of Formula 103 is coupled to a compound of Formula 104. A variety of reaction conditions can be used to effect this coupling, e.g., Sonogashira coupling conditions using a palladium catalyst such as (diphenylphosphineferrocenyl)dichloropalladium or tris (dibenzylidenacetone)-dipalladium; a base such as cesium carbonate or triethylamine; and ligands such as triphenylarsine or triphenylphosphine; Stille conditions using a palladium catalyst such as tris (dibenzylidenacetone)-dipalladium; lithium chloride; and triphenylarsine; or Suzuki coupling conditions wherein the compound of Formula 104 is treated with a borane such as catechol borane or 9-borabicyclo[3.3.1]nonane and the two fragments are coupled using a palladium catalyst such as those described above, a base such as cesium carbonate, and triphenylarsine.

[0073] According to one embodiment, a mixture of a compound of Formula 103; a slight excess (preferably about 1.1 equivalents) of an acetylenic compound of Formula 104; a palladium catalyst such as dichlorobis(triphenylphosphine)palladium(II) and preferably about 0.025 equivalents

of the catalyst; and cuprous iodide in a base such as triethylamine is heated at about 50°C. The corresponding, optionally substituted compound of Formula 105 is isolated and purified.

Preparation of Compounds of Formula 107

[0074] Referring to Reaction Scheme 1, Step 3, the protecting group is then removed from the carboxylic acid. When the protecting group is a lower alkyl ester, it can be removed by treatment with aqueous alcoholic base at an elevated temperature. For example, methyl esters can be converted to the corresponding carboxylic acid by treatment of the ester with potassium hydroxide in a 1:1:1 solution of methanol, THF, and alcohol at about 50°C for 30 minutes. The corresponding, optionally substituted compound of Formula 107 is isolated and used without further purification.

Preparation of Compounds of Formula 109

[0075] Referring to Reaction Scheme 1, Step 4, bis(acetonitrile)dichloropalladium (about 0.050 equivalents) is added to a solution of a compound of Formula 107 and a base, such as triethylamine, in a nonpolar, aprotic solvent such as THF. The solution is maintained at an elevated temperature, preferably about 50°C. The corresponding, optionally substituted isochromen-1-one of Formula 109 is isolated and purified.

Preparation of Compounds of Formula 111

[0076] Referring to Reaction Scheme 1, Step 5, a solution of isochromen-1-one of Formula 109 and an excess (preferably about 3 equivalents) of a primary amine of formula R_1NH_2 in a nonpolar solvent such as toluene is heated at reflux, preferably about 140°C. The resulting amide is isolated and dissolved in a polar, protic solvent such as methanol. Aqueous acid, preferably about 5% aqueous hydrochloric acid is added to the solution which is then heated to about 50°C. Additional aqueous alcoholic acid may be added if required to complete the reaction. The corresponding, optionally substituted 2H-isoquinolin-1-one of Formula 111 is isolated and purified.


Preparation of Compounds of Formula 113


[0077] Referring to Reaction Scheme 1, Step 6, the protecting group is removed from the primary amine. In a preferred embodiment, the protecting group is Boc and its removal can be accomplished through treatment with aqueous TFA at room temperature. The product, a compound of Formula 113 is isolated and can be used without further purification.

[0078] In certain compounds of the invention, particular stereoconfiguration can be preferred for the R₂ substituent, such as the (R) isomer, which can be obtained. An amine of Formula 113 is dissolved in an inert organic solvent (such as IPA) and warmed to 60°C. In a separate vessel, a resolving agent (such as dibenzoyl-D-tartaric acid) is dissolved, preferably in the same warm solvent, and then quickly added (with agitation) to the warm amine solution. The reaction mixture is left to crystallize by cooling to room temperature over 16 hours under continuing agitation. The desired isomer, e.g., the (R) isomer is isolated and purified in the usual manner.

[0079] For the sake of brevity in the remaining description of the synthesis of compounds of Formula I, it should be understood that either single isomer or a mixture of isomers may be employed to give the corresponding product.

Reaction Scheme 2





Preparation of Formula 203

[0080] Referring to Reaction Scheme 2, Step 1, to a solution of a compound of Formula 113 is added successively a slight excess (preferably about 1.2 equivalents) of an aldehyde comprising R_4' (i.e., a compound having the formula $R_4'CHO$ where $R_4'CH_2-$ is equivalent to R_4 and R_4 is as described above or is a protected precursor to such a substituent, e.g., (3-oxo-propyl)-carbamic acid *tert*-butyl ester) and a reducing agent such as sodium triacetoxyborohydride. The resulting mixture is stirred for several hours. The product, a compound of Formula 203 is isolated and purified.

Preparation of Formula 205


[0081] Referring to Reaction Scheme 2, Step 2, to a solution of a compound of Formula 203 and an amine base such as diisopropylethylamine in a nonpolar, aprotic solvent such as dichloromethane is added an R_3 acyl chloride (such as $Cl-C(O)-R_3$ where R_3 is as described above). The resulting solution is stirred under nitrogen at

room temperature for several hours. The product, a compound of Formula 205 is isolated and purified.

Preparation of Formula 207


[0082] Optionally, any protecting groups on compounds of Formula 205 are then removed. For example, if R_4 comprises a protected amine wherein the protecting group is a Boc group, the Boc group can be removed by treatment of the compound of Formula 205 with an acid such as trifluoroacetic acid in a nonpolar, aprotic solvent such as dichloromethane, while maintaining the reaction at about room temperature. The reaction is monitored e.g., by TLC. Upon completion, the product, a compound of Formula 207 is isolated and purified.

Reaction Scheme 3




[0083] Referring to Reaction Scheme 3, to a solution of a compound of Formula 203 and an amine base such as diisopropylethylamine in a nonpolar, aprotic solvent such as dichloromethane is added a compound having the formula $\text{Cl-S(O)}_2\text{-R}_{3a}$ or $\text{O-(S(O)}_2\text{-R}_{3a})_2$ where R_{3a} is as described above. The resulting solution is stirred under nitrogen at room temperature for several hours. The product, a compound of Formula 303 is isolated and purified.


Reaction Scheme 4



[0084] Referring to Reaction Scheme 4, to a solution of a compound of Formula 203 and an amine base such as diisopropylethylamine in a nonpolar, aprotic solvent such as dichloromethane is added a compound having the formula Cl-CH₂-R_{3b} where R_{3b} is as described above. The resulting solution is stirred under nitrogen at room temperature or with heat for several hours. The product, a compound of Formula 403 is isolated and purified.

Reaction Scheme 5





Preparation of Formula 503

[0085] Referring to Reaction Scheme 5, Step 1, to an optionally substituted compound of Formula 113 dissolved in a polar, aprotic solvent (such as DMF) in the presence of a base (such as potassium carbonate) is added one equivalent of an optionally substituted suitably protected aldehyde wherein such aldehyde further comprises a leaving group, preferably, a halide. The solution is heated at reflux, monitoring completion of the reaction (e.g., by TLC). The reaction mixture is cooled and the corresponding, optionally substituted compound of Formula 503 is isolated and purified.

Preparation of Formula 505


[0086] Referring to Reaction Scheme 5, Step 2, to an optionally substituted

compound of Formula 503 in an inert solvent (such as dichloromethane) in the presence of about 1.5 molar equivalents of an amine base (such as triethylamine) is added about 1.5 molar equivalents of an R_9 acid chloride, such as, $Cl-C(O)-R_9$, where R_9 is as described above. The reaction takes place, with stirring, at room temperature over a period of 4 to 24 hours. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 505 is isolated and purified.

Preparation of Formula 507


[0087] Referring to Reaction Scheme 5, Step 3, a solution of a compound of Formula 505 and an excess of ammonium acetate in acetic acid is heated at reflux for 1-4 hours. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 507 is isolated and purified.

Reaction Scheme 6




603

Step 2




605

Step 3



607

Step 4



Preparation of Formula 603

[0088] Referring to Reaction Scheme 6, Step 1, a suspension of a compound of Formula 113, an alpha-haloketone reagent of the Formula $R_{13}'(CO)CH_2X$ wherein X is a leaving group (such as a halide) and R_{13}' is as described herein, and about an equivalent of a base, such as potassium carbonate in a polar, aprotic solvent such as DMF is stirred at room temperature. The reaction is diluted with water and the resulting solid, a compound of Formula 603, is used in the subsequent step without further purification.

Preparation of Formula 605

[0089] Referring to Reaction Scheme 6, Step 2, a solution of the compound of Formula 603, about an equivalent of an amine base, such as triethylamine and about an equivalent of an acid chloride (such as a compound of Formula R_9-COCl) in an organic solvent such as methylene chloride is stirred at room temperature for several hours. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 605 is isolated and purified.


Preparation of Formula 607


[0090] Referring to Reaction Scheme 6, Step 3, a solution of a compound of Formula 605 and an excess of ammonium acetate in acetic acid is heated at reflux using a Dean-Stark trap and condenser. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 607 is isolated and purified.

Preparation of Formula 609

[0091] Optionally, in the event that group R_{13}' comprises a functionality bearing a protecting group, the protecting group is removed. Thus, if R_{13}' further comprises an amine bearing a Pht group, the protecting group is removed as shown in Reaction Scheme 6, Step 4. A solution of a compound of Formula 607 and an excess of anhydrous hydrazine in a polar, protic solvent such as ethanol is heated at reflux. The reaction is cooled to about 5°C and any precipitate is filtered off. The filtrate is concentrated in vacuo and purified to yield a compound of Formula 609.

Reaction Scheme 7





Preparation of Formula 703

[0092] Referring to Reaction Scheme 7, Step 1, reductive amination of amines of Formula 113 (prepared as described in WO 0130768) with an optionally substituted, aldehyde-containing carbamic acid ester (Seki *et. al. Chem. Pharm. Bull.* 1996, 44, 2061) gives urethane intermediates. Removal of the Boc protecting group furnishes an amine of Formula 705.

[0093] More specifically, to a solution of a compound of Formula 113 and an equivalent of a suitably protected aldehyde (Seki *et. al. Chem. Pharm. Bull.* 1996, 44, 2061) in dichloromethane is added a slight excess of a reducing agent, such as sodium triacetoxyborohydride. The resultant cloudy mixture is maintained at ambient

temperature. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 703 is isolated and used in the subsequent step without purification.

Preparation of Formula 705

[0094] Referring to Reaction Scheme 7, Step 2, the amine protecting group, PG, is then removed. For example, when PG is Boc, to a solution of a compound of Formula 703 in a nonpolar, aprotic solvent such as dichloromethane is added a strong acid such as trifluoroacetic acid. The resultant solution is maintained at ambient temperature overnight and concentrated under reduced pressure. The residue is isolated to give a compound of Formula 705 which was used in the subsequent step without purification.


Preparation of Formula 707

[0095] Referring to Reaction Scheme 7, Step 3, to a solution of a compound of Formula 705 in a nonpolar, aprotic solvent such as dichloromethane is added an excess, preferably about two equivalents of an amine base such as triethylamine, followed by about an equivalent or slight excess of an acid chloride of the formula $R_{14}COCl$. The resultant solution is stirred at ambient temperature for about 3 hours. Completion is monitored, e.g., by TLC. The corresponding compound of Formula 707 is isolated and purified.

Preparation of Formula 709

[0096] Referring to Reaction Scheme 7, Step 4, a solution of a compound of Formula 707 in an excess of phosphorus oxychloride is heated at reflux. After 8 hours, the reaction mixture is allowed to cool to ambient temperature and concentrated under reduced pressure. The corresponding compound of Formula 709 is isolated and purified.

Reaction Scheme 8




Preparation of Formula 709

[0097] As an alternative to Steps 3 and 4 of Reaction Scheme 7, acylation of primary amines of Formula 705, followed by acetic acid mediated cyclization, can proceed without isolation of the intermediate amides to provide the target compound of Formula 709. This route is shown in Reaction Scheme 8.


[0098] More specifically, to a solution of a compound of Formula 705 in a nonpolar, aprotic solvent such as dichloromethane is added an excess, preferably about two equivalents of an amine base, such as triethylamine, followed by about an equivalent of an acid chloride of formula $R_{14}COCl$. The resultant solution is stirred at ambient temperature for 2 hours, then evaporated under reduced pressure. The resultant solid is treated with glacial acetic acid, then the resultant suspension is heated at reflux for about 48 hours. The reaction is cooled to ambient temperature then evaporated under reduced pressure. The corresponding compound of Formula 709 is isolated and purified.

Reaction Scheme 9




[0099] Referring to Reaction Scheme 9, a compound of Formula 203 is reacted with a slight excess of a compound of the formula $R_{15}O(CO)Cl$ in the presence of a base such as triethylamine in a nonpolar, aprotic solvent such as dichloromethane. The product, a compound of Formula 903 is isolated and purified.

Reaction Scheme 10



[00100] Referring to Reaction Scheme 10, a compound of Formula 203 is treated with a slight excess of an isocyanate $R_{17}-N=C=O$ in the presence of a base, such as triethylamine, in a nonpolar, aprotic solvent, such as dichloromethane. The product, a compound of Formula 1003, is isolated and purified.


Reaction Scheme 11




Preparation of Compounds of Formula 1103

[00101] Referring to Reaction Scheme 11, to a solution of a compound of Formula 1101 in a nonpolar, aprotic solvent such as DMF are added a base such as triethylamine and an excess (preferably, about 1.5 equivalents) of imidazole followed by about an equivalent of tetrabutylammonium iodide. The resultant solution is heated to about 90 °C, stirred for about 18 h and allowed to cool to room temperature. The product, a compound of Formula 1103, is isolated and purified.

Reaction Scheme 12





Preparation of Compounds of Formula 1203

[00102] Referring to Reaction Scheme 12, Step 1, to a solution of a compound of Formula 113 and an excess of an optionally substituted, aldehyde-containing carbamic acid ester such as ((*S*)-4-benzyloxycarbonylamino-1-formyl-butyl)-carbamic acid *tert*-butyl ester) in a nonpolar, aprotic solvent such as CH_2Cl_2 is added sodium triacetoxyborohydride. The mixture is stirred overnight. The product, a compound of Formula 1203, is isolated and purified.

Preparation of Compounds of Formula 1205

[00103] Referring to Reaction Scheme 12, Step 2, to a solution of a compound


of Formula 1203 in a nonpolar, aprotic solvent such as toluene is added a base such as triethylamine followed by dropwise addition of an excess of an acid chloride of the formula $R_{14}-COCl$. The reaction mixture is heated to about 80 °C for about 18 h, then at reflux for about 4 h. The product, a compound of Formula 1205, is isolated and purified.

Preparation of Compounds of Formula 1207

[00104] Referring to Reaction Scheme 12, Step 3, a solution of a compound of Formula 1205 in a solvent such as CH_2Cl_2 /TFA (preferably, about 4:1 CH_2Cl_2 /TFA) is stirred at room temperature. The reaction mixture is concentrated under reduced pressure and the residue is diluted with a nonpolar, aprotic solvent such as CH_2Cl_2 and washed with aqueous base. The aqueous layer is extracted with a nonpolar, aprotic solvent such as CH_2Cl_2 and the combined extracts are dried, filtered and concentrated under reduced pressure. The residue is diluted with a nonpolar, aprotic solvent such as THF and aqueous base (preferably, saturated aqueous $NaHCO_3$). The mixture is stirred at room temperature for 10 days. The product, a compound of Formula 1207, is isolated and purified.

Particular Processes and Last Steps

[00105] A compound of Formula 203:



(where R_2 is optionally protected) is contacted with a slight molar excess of an R_3 chloride [such as, $Cl-C(O)-R_3$, $Cl-S(O)_2-R_{3a}$, $Cl-CH_2-R_{3b}$, $Cl-C(O)-O-R_{15}$ and $Cl-S(O)_2-NH-R_{3a}$] or an isocyanate (such as $O=C=N-R_{17}$) or an anhydride (such as $O[C(O)R_{15}]_2$ or $O[S(O)_2R_{3a}]_2$) to give the corresponding optionally protected compound

of Formula I.

[00106] A compound of Formula 505, 605, 705, or 707 is optionally cyclized by acid-mediated cyclization.

[00107] A racemic mixture of isomers of a compound of Formula I is optionally placed on a chromatography column and separated into (R)- and (S)-enantiomers.

[00108] A compound of Formula I is optionally contacted with a pharmaceutically acceptable acid or base to form the corresponding acid or base addition salt.

[00109] A pharmaceutically acceptable acid addition salt of a compound of Formula I is optionally contacted with a base to form the corresponding free base of Formula I.

[00110] A pharmaceutically acceptable base addition salt of a compound of Formula I is optionally contacted with an acid to form the corresponding free acid of Formula I.

Particular Embodiments of Compounds of the Invention

R₁

[00111] When considering the compounds of Formula I, in one embodiment, R₁ is selected from hydrogen, optionally substituted C₁-C₈ alkyl-, optionally substituted aryl-, optionally substituted heteroaryl-, optionally substituted aryl-C₁-C₄-alkyl-, and optionally substituted heteroaryl-C₁-C₄-alkyl- (more particularly optionally substituted aryl and optionally substituted aryl-C₁-C₄-alkyl-). In a more particular embodiment R₁ is selected from hydrogen, optionally substituted C₁-C₄ alkyl-, optionally substituted phenyl-C₁-C₄-alkyl-, optionally substituted naphthalenylmethyl-, optionally substituted phenyl-, and naphthyl-. Even more particularly, R₁ is optionally substituted phenyl-C₁-C₄-alkyl-, optionally substituted heteroaryl-C₁-C₄-alkyl-, or naphthalenylmethyl-.

[00112] Yet more particularly, R₁ is naphthyl-, phenyl-, bromophenyl-, chlorophenyl-, methoxyphenyl-, ethoxyphenyl-, tolyl-, dimethylphenyl-, chorofluorophenyl-, methylchlorophenyl-, ethylphenyl-, phenethyl-, benzyl-, chlorobenzyl-, methylbenzyl-, methoxybenzyl-, cyanobenzyl-, hydroxybenzyl-, dichlorobenzyl-, dimethoxybenzyl-, or naphthalenylmethyl-. More suitably, R₁ is

benzyl-, cyanobenzyl-, methoxybenzyl-, or naphthalenylmethyl-. Most particularly, R_1 is benzyl-.

R_2 and $R_{2'}$

[00113] When considering the compounds of Formula I and as will be appreciated by those skilled in the art, the compounds described herein possess a potentially chiral center at the carbon to which R_2 and $R_{2'}$ are attached. The R_2 and $R_{2'}$ groups may be the same or different; if different, the compound is chiral (i.e., has a stereogenic center). When R_2 and $R_{2'}$ are different, in particular embodiments $R_{2'}$ is hydrogen and R_2 is other than hydrogen. The invention contemplates the use of pure enantiomers and mixtures of enantiomers, including racemic mixtures, although the use of a substantially optically pure enantiomer will generally be preferred. The term "substantially pure" means having at least about 95% chemical purity with no single impurity greater than about 1%. The term "substantially optically pure" or "enantiomerically pure" means having at least about 97.5% enantiomeric excess. In a particular embodiment, the stereogenic center to which R_2 and $R_{2'}$ are attached is of the R configuration.

[00114] When considering the compounds of Formula I, in one embodiment, R_2 and $R_{2'}$ are independently chosen from hydrogen, optionally substituted alkyl-, optionally substituted alkoxy, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-.

[00115] In another embodiment, R_2 and $R_{2'}$ taken together form a 3- to 7-membered ring which may optionally be substituted one or more of the following groups: hydroxyl, halogen (particularly chloro and fluoro), optionally substituted C_1 - C_4 alkyl- (particularly methyl-), C_1 - C_4 alkoxy (particularly methoxy), cyano, amino, substituted amino, or carbamyl-.

[00116] In one embodiment, R_2 is optionally substituted C_1 - C_4 alkyl-, and $R_{2'}$ is hydrogen or optionally substituted C_1 - C_4 alkyl-. More suitably, R_2 is hydrogen and $R_{2'}$ is optionally substituted C_1 - C_4 alkyl-. In a most particular embodiment R_2 is chosen from methyl-, ethyl-, propyl (particularly, *c*-propyl or *i*-propyl), butyl (particularly, *t*-butyl), methylthioethyl-, methylthiomethyl-, aminobutyl-, (CBZ)aminobutyl-, cyclohexylmethyl-, benzyloxymethyl-, methylsulfanylethyl-, methylsulfanylmethyl-, and hydroxymethyl-, and $R_{2'}$ is hydrogen. Especially chosen

embodiments are when R_2' is hydrogen and R_2 is ethyl or propyl (particularly, *c*-propyl or *i*-propyl). Even more particularly, R_2 is *i*-propyl. Yet more particularly, the stereogenic center to which R_2 and R_2' is attached is of the *R* configuration.


[00117] In one embodiment, if either R_2 or R_2' is hydrogen, then the other is not hydrogen. In another embodiment, both R_2 and R_2' are hydrogen.

R_5 , R_6 , R_7 , and R_8

[00118] In other embodiments R_5 , R_6 , R_7 , and R_8 are independently chosen from hydrogen, hydroxyl, halogen (particularly chloro and fluoro), optionally substituted C_1 - C_4 alkyl- (particularly methyl-), C_1 - C_4 alkoxy (particularly methoxy), cyano, amino, substituted amino, or carbamyl-. More particularly, R_5 , R_6 , R_7 , and R_8 are methoxy, methyl, trifluoromethyl, cyano, hydrogen or halo. In particular embodiments, R_5 is hydrogen or halo; R_6 is hydrogen, optionally substituted C_1 - C_4 alkyl- (particularly, methyl-) or halo; R_7 is hydrogen, halo, optionally substituted C_1 - C_4 alkyl- (particularly, methyl- or trifluoromethyl-), C_1 - C_4 alkoxy (particularly, methoxy), cyano, substituted amino, or carbamyl-; and R_8 is hydrogen, C_1 - C_4 alkyl- (particularly, methyl-), C_1 - C_4 alkoxy (particularly, methoxy), hydroxy, or halo. Still more particularly are the compounds where only one of R_5 , R_6 , R_7 , and R_8 is not hydrogen, especially R_7 . Another particular embodiment is drawn to the compounds where R_7 and R_8 are not hydrogen.

Compounds wherein R_{12} is an Optionally Substituted Imidazolyl

[00119] When R_{12} is an optionally substituted imidazolyl-, in particular embodiments, R_{12} has the formula:



wherein

R_9 is chosen from hydrogen, optionally substituted C_1 - C_8 alkyl-, optionally substituted aryl-, optionally substituted aryl- C_1 - C_4 -alkyl-, optionally substituted heteroaryl- C_1 - C_4 -alkyl-, optionally substituted aryl- C_1 - C_4 -alkoxy, optionally substituted heteroaryl- C_1 - C_4 -alkoxy, and optionally substituted heteroaryl-; and


R_{13} and $R_{13'}$ are independently hydrogen, optionally substituted C_1 - C_8 alkyl-, optionally substituted aryl-, or optionally substituted aryl- C_1 - C_4 -alkyl-.

[00120] According to one embodiment, R_9 is lower-alkyl; phenyl substituted with optionally substituted C_1 - C_4 -alkyl-, C_1 - C_4 -alkoxy-, and/or halo; phenyl-; benzyl-; thiophenyl-; or thiophenyl- substituted with C_1 - C_4 -alkyl-, C_1 - C_4 -alkoxy-, and/or halo. More suitably, R_9 is lower-alkyl; phenyl-; or phenyl- substituted with one or more of the following groups: methyl, methoxy, trifluoromethyl, or halo.

[00121] According to another embodiment, R_{13} is hydrogen and $R_{13'}$ is substituted C_1 - C_4 alkyl-. More suitably, R_{13} is hydrogen and $R_{13'}$ is aminomethyl-, aminoethyl-, aminopropyl-, acetylamino-methyl-, acetylaminoethyl-, benzyloxycarbonylamino-methyl- or benzyloxycarbonylamino-ethyl-.

Compounds wherein R_{12} is an Optionally Substituted Imidazoliny

[00122] When R_{12} is an optionally substituted imidazoliny, in one embodiment, R_{12} has the formula



wherein,

R_{14} is chosen from hydrogen, optionally substituted C_1 - C_8 alkyl-, optionally substituted aryl-, optionally substituted aryl- C_1 - C_4 -alkyl-, optionally substituted heteroaryl-, optionally substituted heteroaryl- C_1 - C_4 -alkyl-; and

R_{10} , $R_{10'}$, R_{11} and $R_{11'}$ are independently chosen from hydrogen, optionally substituted C_1 - C_8 alkyl-, optionally substituted aryl-, and optionally substituted aryl- C_1 - C_4 -alkyl-.

[00123] In one embodiment, R_{14} is lower-alkyl-; methylenedioxyphenyl-;

phenyl-; phenyl substituted with optionally substituted C₁-C₄ alkyl-, C₁-C₄ alkoxy-, and/or halo; benzyl-; thienyl substituted with C₁-C₄ alkyl; benzyl; thiophenyl-; or thiophenyl- substituted with C₁-C₄-alkyl-, C₁-C₄-alkoxy-, and/or halo. More suitably, R₁₄ is lower-alkyl; phenyl-; or phenyl- substituted with one or more of the following groups: methyl, methoxy, trifluoromethyl, or halo.

[00124] In one embodiment, R₁₀, R_{10'}, R_{11'}, and R₁₁ are independently hydrogen or optionally substituted C₁-C₄ alkyl-. More suitably, R_{11'} and R₁₁ are hydrogen.

Compounds wherein R₁₂ is -NHR₄, -NR₄(COR₃), NR₄(SO₂R_{3a}), or -NR₄(CH₂R_{3b})
R₄

[00125] In one embodiment, R₄ is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, and optionally substituted heterocyclyl-. More suitably, R₄ is R₁₆-alkylene-, and R₁₆ is chosen from alkoxy, amino, alkylamino, dialkylamino, carboxy, guanidine, hydroxyl-, and N-heterocyclyl-.

[00126] In a more particular embodiment, R₄ is selected from optionally substituted lower-alkyl-, optionally substituted cyclohexyl-; phenyl substituted with hydroxy, lower-alkoxy or lower-alkyl-; benzyl-; heteroarylmethyl-; heteroarylethyl-; and heteroarylpropyl-.

[00127] In a most particular embodiment, R₄ is chosen from methyl-, ethyl-, propyl-, butyl-, cyclohexyl-, carboxyethyl-, carboxymethyl-, methoxyethyl-, hydroxyethyl-, hydroxypropyl-, dimethylaminoethyl-, dimethylaminopropyl-, diethylaminoethyl-, diethylaminopropyl-, aminopropyl-, methylaminopropyl-, 2,2-dimethyl-3-(dimethylamino)propyl-, 1-cyclohexyl-4-(diethylamino)butyl-, aminoethyl-, aminobutyl-, aminopentyl-, aminohexyl-, aminoethoxyethyl-, isopropylaminopropyl-, diisopropylaminoethyl-, 1-methyl-4-(diethylamino)butyl-, (t-Boc)aminopropyl-, hydroxyphenyl-, benzyl-, methoxyphenyl-, methylmethoxyphenyl-, dimethylphenyl-, tolyl-, ethylphenyl-, (oxopyrrolidinyl)propyl-, (methoxycarbonyl)ethyl-, benzylpiperidinyl-, pyridinylethyl-, pyridinylmethyl-, morpholinylethyl morpholinylpropyl-, piperidinyl-, azetidinylmethyl-, azetidinylethyl-, azetidinypropyl-, pyrrolidinylmethyl-, pyrrolidinylethyl-, pyrrolidinylpropyl-, piperidinylmethyl-, piperidinylethyl-, imidazolylpropyl-, imidazolylethyl-, (ethylpyrrolidinyl)methyl-, (methylpyrrolidinyl)ethyl-, (methylpiperidinyl)propyl-,

(methylpiperazinyl)propyl-, guanidino-methyl-, guanidino-ethyl-, guanidino-propyl-, furanylmethyl and indolyethyl- (most especially aminopropyl).

R₃

[00128] When considering the compounds of Formula I, in a particular embodiment R₃ is selected from hydrogen, optionally substituted alkyl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, optionally substituted heteroaryl-, optionally substituted aryl-, R₁₅O- and R₁₇-NH-, wherein R₁₅ is chosen from optionally substituted alkyl and optionally substituted aryl and R₁₇ is chosen from hydrogen, optionally substituted alkyl and optionally substituted aryl-.

[00129] In a more particular embodiment, when R₃ is not R₁₇NH- or R₁₅O-, R₃ is chosen from optionally substituted alkyl-, aryl- (including phenyl-, biphenyl-, and naphthyl-); substituted aryl- (including phenyl substituted with one or more cyano, halo, lower-alkyl-, lower-alkoxy, hydroxy-loweralkyl-, nitro, carboxy, methylenedioxy, trifluoromethoxy, or trifluoromethyl-); benzyl-; and optionally substituted heteroaryl-.

[00130] In a most particular embodiment, when R₃ is not R₁₇NH- or R₁₅O-, R₃ is chosen from ethyl-, propyl-, chloropropyl-, butoxy, heptyl-, butyl-, octyl-, tridecanyl-, (ethoxycarbonyl)ethyl-, dimethylaminoethyl-, dimethylaminomethyl-, phenyl-, naphthyl-, halophenyl-, polyhalophenyl-, cyanophenyl-, hydroxymethylphenyl-, halo(trifluoromethyl)phenyl-, chlorophenoxymethyl-, methoxyphenyl-, carboxyphenyl-, ethylphenyl-, tolyl-, hydroxymethylphenyl-, ethylphenyl-, biphenyl-, methylenedioxyphenyl-, methylsulfonylphenyl-, methoxychlorophenyl-, chloronaphthyl-, acetylphenyl-, methylhalophenyl-, trifluoromethylphenyl-, trifluoromethoxyphenyl-, butylphenyl-, pentylphenyl-, methylnitrophenyl-, phenoxymethyl-, dimethoxyphenyl-, phenylvinyl-, nitrochlorophenyl-, nitrophenyl-, dinitrophenyl-, bis(trifluoromethyl)phenyl-, benzyloxymethyl-, benzyl-, furanyl-, benzofuranyl-, pyridinyl-, pyridyl-, indolyl-, methylpyridinyl-, methylpyridyl-, (3-carbamoyl)pyridinyl- [nicotinamide], 3-carbamoyl-6-methylpyridinyl-, quinolinyl-, picolinyl-, pyrazolyl-, pyrazinyl-, methylpyrazinyl-, morpholinomethyl-, methylthiomethyl-, methoxymethyl-, imidazolyl-, isoxazolyl-, methyl-isoxazolyl-, benzothiadiazolyl-, methylenedioxyphenyl-, thienyl-, methylthienyl-, methyl-nicotinamidyl-, methyl-

pyrazinyl; benzodioxolyl; and methyl-thiophenyl-.

[00131] More suitably, R_3 is tolyl-, halophenyl-, halomethylphenyl-, hydroxymethylphenyl-, methylenedioxyphenyl-, formylphenyl or cyanophenyl-.

[00132] In another particular embodiment, when R_3 is $R_{17}NH$ -, R_{17} is chosen from lower-alkyl-, cyclohexyl-, phenyl-, and phenyl substituted with halo, lower-alkyl-, loweralkoxy, or lower-alkylsulfanyl-.

[00133] In another particular embodiment, when R_3 is $R_{17}NH$ -, R_{17} is isopropyl-, butyl-, cyclohexyl-, phenyl-, bromophenyl-, dichlorophenyl-, methoxyphenyl-, ethylphenyl-, tolyl-, trifluoromethylphenyl or methylthiophenyl-.

[00134] In a particular embodiment, when R_3 is $R_{15}O$ -, R_{15} is chosen from lower-alkyl-, cyclohexyl-, phenyl-, and phenyl substituted with halo, lower-alkyl-, loweralkoxy, or lower-alkylsulfanyl-.

[00135] In a most particular embodiment, when R_3 is $R_{15}O$ -, R_{15} is isopropyl-, butyl-, cyclohexyl-, phenyl-, bromophenyl-, dichlorophenyl-, methoxyphenyl-, ethylphenyl-, tolyl-, trifluoromethylphenyl or methylthiophenyl-.

R_{3a}

[00136] When R_{12} is $-NR_4(SO_2R_{3a})$, R_4 is as described above and R_{3a} is chosen from C_1 - C_{13} alkyl-, phenyl-, naphthyl-, phenyl substituted with cyano, halo, lower-alkyl-, lower-alkoxy, nitro, methylenedioxy, or trifluoromethyl-, biphenyl and heteroaryl-. More suitably, R_{3a} is chosen from phenyl substituted with halo, lower-alkyl-, lower-alkoxy, cyano, nitro, methylenedioxy, or trifluoromethyl-, and naphthyl-.

R_{3b}

[00137] When R_{12} is $-NR_4(CH_2R_{3b})$, R_4 is as described above and R_{3b} is chosen from C_1 - C_{13} alkyl-, substituted lower-alkyl-, phenyl-, naphthyl-, phenyl substituted with cyano, halo, lower-alkyl-, lower-alkoxy, nitro, methylenedioxy, or trifluoromethyl-, biphenyl-, benzyl and heterocyclyl-. Most suitably, R_{3b} is chosen from phenyl substituted with one or more halo, methyl-, cyano, trifluoromethyl-, trifluoromethoxy, carboxy, or methoxycarbonyl groups; piperidinyl-, and naphthyl-. Even more suitably, R_{3b} is halophenyl-, methylhalophenyl-, polyhalophenyl-, tolyl-, dimethylphenyl-, methoxyphenyl-, dimethoxyphenyl-, cyanophenyl-, trifluoromethylphenyl-, trifluoromethoxyphenyl-, bis(trifluoromethyl)phenyl-,

carboxyphenyl-, t-butylphenyl-, methoxycarbonylphenyl-, piperidinyl-, and naphthyl-.

Particular Subgenus

[00138] In a particular subgenus of compounds of Formula I,

R_1 is optionally substituted aryl- C_1 - C_4 -alkyl-, optionally substituted heteroaryl- C_1 - C_4 -alkyl-, or naphthalenylmethyl;

R_2 is optionally substituted C_1 - C_4 -alkyl-;

R_2' is hydrogen;

R_7 is hydrogen, methyl, or cyano;

R_5 , R_6 , and R_8 are hydrogen;

and

R_{12} is optionally substituted imidazolyl-, optionally substituted imidazolyl-, $-NHR_4$; $-N(R_4)(COR_3)$; $-N(R_4)(SO_2 R_{3a})$; or $-N(R_4)(CH_2 R_{3b})$.

[00139] In a particular subgenus of compounds of Formula I wherein R_{12} is optionally substituted imidazolyl-,

R_1 , R_2 , R_2' , R_5 , R_6 , R_7 , and R_8 are as defined above;

R_9 is lower-alkyl; phenyl-; or phenyl- substituted with one or more of the following groups: methyl, methoxy, trifluoromethyl, or halo; and

R_{13} is hydrogen and $R_{13'}$ is substituted C_1 - C_4 alkyl- (especially, aminomethyl-, aminoethyl-, aminopropyl-, acetylaminomethyl-, acetylaminethyl-, benzyloxycarbonylamino-methyl- or benzyloxycarbonylamino-ethyl-.)

[00140] In a particular subgenus of compounds of Formula I wherein R_{12} is optionally substituted imidazolyl-,

R_1 , R_2 , R_2' , R_5 , R_6 , R_7 , and R_8 are as defined above;

R_{14} is lower-alkyl; phenyl-; or phenyl- substituted with one or more of the following groups: methyl, methoxy, trifluoromethyl, or halo; and

R_{10} , $R_{10'}$, $R_{11'}$, and R_{11} are independently hydrogen or optionally substituted C_1 - C_4 alkyl-. More particularly, $R_{11'}$ and R_{11} are hydrogen.

[00141] In a particular subgenus of compounds of Formula I wherein R_{12} is $-NHR_4$,

R_1 , R_2 , R_2' , R_5 , R_6 , R_7 , and R_8 are as defined above;

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, and

optionally substituted heterocyclyl-.

[00142] In a particular subgenus of compounds of Formula I wherein R_{12} is –
 $NR_4(COR_3)$,

$R_1, R_2, R_2', R_5, R_6, R_7$, and R_8 are as defined above;

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, and optionally substituted heterocyclyl- and

R_3 is selected from hydrogen, optionally substituted alkyl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, optionally substituted heteroaryl-, optionally substituted aryl-, $R_{15}O-$ and $R_{17}NH-$, wherein R_{15} is chosen from optionally substituted alkyl and optionally substituted aryl and R_{17} is chosen from hydrogen, optionally substituted alkyl and optionally substituted aryl.

[00143] More particularly, $R_1, R_2, R_2', R_5, R_6, R_7$, and R_8 are as defined above;

R_4 is selected from optionally substituted lower-alkyl-, optionally substituted cyclohexyl-, phenyl substituted with hydroxy, lower-alkoxy, or lower-alkyl-, benzyl-, heteroarylmethyl-, heteroarylethyl-, heteroarylpropyl- (especially R_4 is R_{16} -alkylene-, and R_{16} is chosen from alkoxy, amino, alkylamino, dialkylamino, carboxy, hydroxyl-, and N-heterocyclyl-);

and

R_3 is selected from hydrogen, optionally substituted alkyl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, optionally substituted heteroaryl-, optionally substituted aryl-, $R_{15}O-$ and $R_{17}NH-$, wherein R_{15} is chosen from optionally substituted alkyl and optionally substituted aryl and R_{17} is chosen from hydrogen, optionally substituted alkyl and optionally substituted aryl.

[00144] In a particular subgenus of compounds of Formula I wherein R_{12} is –
 $NR_4(COR_3)$,

$R_1, R_2, R_2', R_5, R_6, R_7$, and R_8 are as defined above;

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, and optionally substituted heterocyclyl- and

and

R_3 is selected from optionally substituted alkyl-, aryl-, substituted aryl-, benzyl-, and optionally substituted heteroaryl-.

[00145] In a particular subgenus of compounds of Formula I wherein R_{12} is —
 $NR_4(COR_3)$,

$R_1, R_2, R_2', R_5, R_6, R_7$, and R_8 are as defined above;

R_3 is tolyl-, halophenyl-, halomethylphenyl-, hydroxymethylphenyl-,
 methylenedioxyphenyl-, formylphenyl or cyanophenyl-; and

R_4 is optionally substituted lower-alkyl-, optionally substituted cyclohexyl-;
 phenyl substituted with hydroxy, lower-alkoxy or lower-alkyl-; benzyl-;
 heteroaryl-methyl-; heteroaryl-ethyl-; or heteroaryl-propyl- (especially R_4 is R_{16} -
 alkylene- and R_{16} is alkoxy, amino, alkylamino, dialkylamino, carboxy, guanidine,
 hydroxyl, or N-heterocycl-yl-).

[00146] In a particular subgenus of compounds of Formula I wherein R_{12} is —
 $N(R_4)(CH_2 R_{3b})$,

$R_1, R_2, R_2', R_5, R_6, R_7$, and R_8 are as defined above;

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted
 aryl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, and
 optionally substituted heterocycl-yl- and

R_{3b} is chosen from phenyl substituted with one or more halo, methyl-, cyano,
 trifluoromethyl-, trifluoromethoxy, carboxy, or methoxycarbonyl groups;
 piperidin-yl-; and naphth-yl-.

[00147] In a particular subgenus of compounds of Formula I wherein R_{12} is —
 $NR_4(SO_2 R_{3a})$,

$R_1, R_2, R_2', R_5, R_6, R_7$, and R_8 are as defined above;

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally
 substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-,
 and optionally substituted heterocycl-yl- and

R_{3a} is chosen from phenyl substituted with halo, lower-alkyl-, lower-alkoxy,
 cyano, nitro, methylenedioxy, or trifluoromethyl-; and naphth-yl-.

[00148] Particular compounds include:

N-(3-Amino-propyl)-N-[1-(2-benzyl-4-methoxy-1-oxo-1,2-dihydro-isoquinolin-3-yl)-2-
 methyl-propyl]-4-methyl-benzamide;

N-(3-Amino-propyl)-N-[1-(2-benzyl-4-methoxy-1-oxo-1,2-dihydro-isoquinolin-3-yl)-2-
 methyl-propyl]-2-methoxy-acetamide;

3-(1-Amino-2-methyl-propyl)-2-benzyl-6-chloro-2H-isoquinolin-1-one;

N-(3-Amino-propyl)-N-[1-(2-benzyl-6-chloro-1-oxo-1,2-dihydro-isoquinolin-3-yl)-2-methyl-propyl]-4-methyl-benzamide;
 N-(3-Amino-propyl)-N-[1-(2-benzyl-6-chloro-1-oxo-1,2-dihydro-isoquinolin-3-yl)-2-methyl-propyl]-2-methoxy-acetamide;
 N-(3-Amino-propyl)-N-[1-(2-benzyl-6-chloro-1-oxo-1,2-dihydro-isoquinolin-3-yl)-2-methyl-propyl]-4-bromo-benzamide;
 2-Benzyl-6-chloro-3-[1-(4,4-dimethyl-2-p-tolyl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-2H-isoquinolin-1-one;
 2-Benzyl-6-chloro-3-{1-[2-(3,4-dimethoxy-phenyl)-4,4-dimethyl-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-2H-isoquinolin-1-one;
 2-Benzyl-6-chloro-3-{1-[2-(3,4-dichloro-phenyl)-4,4-dimethyl-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-2H-isoquinolin-1-one;
 2-Benzyl-6-chloro-3-[1-(4,4-dimethyl-2-m-tolyl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-2H-isoquinolin-1-one;
 2-Benzyl-6-chloro-3-{1-[4,4-dimethyl-2-(3-trifluoromethyl-phenyl)-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-2H-isoquinolin-1-one;
 2-Benzyl-6-chloro-3-{1-[2-(3-fluoro-phenyl)-4,4-dimethyl-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-2H-isoquinolin-1-one;
 2-Benzyl-6-chloro-3-{1-[2-(3-fluoro-4-methyl-phenyl)-4,4-dimethyl-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-2H-isoquinolin-1-one;
 2-Benzyl-6-chloro-3-[1-(2-isopropyl-4,4-dimethyl-4,5-dihydro-imidazol-1-yl)-2-methyl-propyl]-2H-isoquinolin-1-one;
 2-Benzyl-6-chloro-3-{1-[2-(4-methoxy-phenyl)-4,4-dimethyl-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-2H-isoquinolin-1-one; and
 2-Benzyl-3-{1-[2-(4-bromo-phenyl)-4,4-dimethyl-4,5-dihydro-imidazol-1-yl]-2-methyl-propyl}-6-chloro-2H-isoquinolin-1-one.

Utility, Testing and Administration

General Utility

[00149] Once made, the compounds of the invention find use in a variety of applications involving alteration of mitosis. As will be appreciated by those skilled in the art, mitosis may be altered in a variety of ways; that is, one can affect mitosis

either by increasing or decreasing the activity of a component in the mitotic pathway. Stated differently, mitosis may be affected (e.g., disrupted) by disturbing equilibrium, either by inhibiting or activating certain components. Similar approaches may be used to alter meiosis.

[00150] In a particular embodiment, the compounds of the invention are used to inhibit mitotic spindle formation, thus causing prolonged cell cycle arrest in mitosis. By "inhibit" in this context is meant decreasing or interfering with mitotic spindle formation or causing mitotic spindle dysfunction. By "mitotic spindle formation" herein is meant organization of microtubules into bipolar structures by mitotic kinesins. By "mitotic spindle dysfunction" herein is meant mitotic arrest and monopolar spindle formation.

[00151] The compounds of the invention are useful to bind to, and/or inhibit the activity of, a mitotic kinesin, KSP. In one embodiment, the KSP is human KSP, although the compounds may be used to bind to or inhibit the activity of KSP kinesins from other organisms. In this context, "inhibit" means either increasing or decreasing spindle pole separation, causing malformation, i.e., splaying, of mitotic spindle poles, or otherwise causing morphological perturbation of the mitotic spindle. Also included within the definition of KSP for these purposes are variants and/or fragments of KSP. See U.S. Patent 6,437,115, hereby incorporated by reference in its entirety. The compounds of the invention have been shown to have specificity for KSP. However, the present invention includes the use of the compounds to bind to or modulate other mitotic kinesins.

[00152] The compounds of the invention are used to treat cellular proliferation diseases. Such disease states which can be treated by the compounds, compositions and methods provided herein include, but are not limited to, cancer (further discussed below), autoimmune disease, fungal disorders, arthritis, graft rejection, inflammatory bowel disease, cellular proliferation induced after medical procedures, including, but not limited to, surgery, angioplasty, and the like. Treatment includes inhibiting cellular proliferation. It is appreciated that in some cases the cells may not be in an abnormal state and still require treatment. Thus, in one embodiment, the invention herein includes application to cells or individuals afflicted or subject to impending affliction with any one of these disorders or states.

[00153] The compounds, compositions and methods provided herein are

particularly deemed useful for the treatment of cancer including solid tumors such as skin, breast, brain, cervical carcinomas, testicular carcinomas, etc. More particularly, cancers that may be treated by the compounds, compositions and methods of the invention include, but are not limited to: Cardiac: sarcoma (angiosarcoma, fibrosarcoma, rhabdomyosarcoma, liposarcoma), myxoma, rhabdomyoma, fibroma, lipoma and teratoma; Lung: bronchogenic carcinoma (squamous cell, undifferentiated small cell, undifferentiated large cell, adenocarcinoma), alveolar (bronchiolar) carcinoma, bronchial adenoma, sarcoma, lymphoma, chondromatous hamartoma, mesothelioma; Gastrointestinal: esophagus (squamous cell carcinoma, adenocarcinoma, leiomyosarcoma, lymphoma), stomach (carcinoma, lymphoma, leiomyosarcoma), pancreas (ductal adenocarcinoma, insulinoma, glucagonoma, gastrinoma, carcinoid tumors, vipoma), small bowel (adenocarcinoma, lymphoma, carcinoid tumors, Kaposi's sarcoma, leiomyoma, hemangioma, lipoma, neurofibroma, fibroma), large bowel (adenocarcinoma, tubular adenoma, villous adenoma, hamartoma, leiomyoma); Genitourinary tract: kidney (adenocarcinoma, Wilm's tumor (nephroblastoma), lymphoma, leukemia), bladder and urethra (squamous cell carcinoma, transitional cell carcinoma, adenocarcinoma), prostate (adenocarcinoma, sarcoma), testis (seminoma, teratoma, embryonal carcinoma, teratocarcinoma, choriocarcinoma, sarcoma, interstitial cell carcinoma, fibroma, fibroadenoma, adenomatoid tumors, lipoma); Liver: hepatoma (hepatocellular carcinoma), cholangiocarcinoma, hepatoblastoma, angiosarcoma, hepatocellular adenoma, hemangioma; Bone: osteogenic sarcoma (osteosarcoma), fibrosarcoma, malignant fibrous histiocyoma, chondrosarcoma, Ewing's sarcoma, malignant lymphoma (reticulum cell sarcoma), multiple myeloma, malignant giant cell tumor chordoma, osteochondroma (osteochondilagenous exostoses), benign chondroma, chondroblastoma, chondromyxofibroma, osteoid osteoma and giant cell tumors; Nervous system: skull (osteoma, hemangioma, granuloma, xanthoma, osteitis deformans), meninges (meningioma, meningiosarcoma, gliomatosis), brain (astrocytoma, medulloblastoma, glioma, ependymoma, germinoma (pinealoma), glioblastoma multiform, oligodendroglioma, schwannoma, retinoblastoma, congenital tumors), spinal cord neurofibroma, meningioma, glioma, sarcoma); Gynecological: uterus (endometrial carcinoma), cervix (cervical carcinoma, pre-tumor cervical dysplasia), ovaries (ovarian carcinoma (serous cystadenocarcinoma, mucinous cystadenocarcinoma,

unclassified carcinoma), granulosa-thecal cell tumors, Sertoli-Leydig cell tumors, dysgerminoma, malignant teratoma), vulva (squamous cell carcinoma, intraepithelial carcinoma, adenocarcinoma, fibrosarcoma, melanoma), vagina (clear cell carcinoma, squamous cell carcinoma, botryoid sarcoma (embryonal rhabdomyosarcoma), fallopian tubes (carcinoma); Hematologic: blood (myeloid leukemia (acute and chronic), acute lymphoblastic leukemia, chronic lymphocytic leukemia, myeloproliferative diseases, multiple myeloma, myelodysplastic syndrome), Hodgkin's disease, non-Hodgkin's lymphoma (malignant lymphoma); Skin: malignant melanoma, basal cell carcinoma, squamous cell carcinoma, Kaposi's sarcoma, moles dysplastic nevi, lipoma, angioma, dermatofibroma, keloids, psoriasis; and Adrenal glands: neuroblastoma. Thus, the term "cancerous cell" as provided herein, includes a cell afflicted by any one of the above identified conditions.

Testing

[00154] For assay of KSP-modulating activity, generally either KSP or a compound according to the invention is non-diffusably bound to an insoluble support having isolated sample receiving areas (e.g., a microtiter plate, an array, etc.). The insoluble support may be made of any material to which the sample can be bound, is readily separated from soluble material, and is otherwise compatible with the overall method of screening. The surface of such supports may be solid or porous and of any convenient shape. Examples of suitable insoluble supports include microtiter plates, arrays, membranes and beads. These are typically made of glass, plastic (e.g., polystyrene), polysaccharides, nylon or nitrocellulose, Teflon™, etc. Microtiter plates and arrays are especially convenient because a large number of assays can be carried out simultaneously, using small amounts of reagents and samples. The particular manner of binding of the sample is not crucial so long as it is compatible with the reagents and overall methods of the invention, maintains the activity of the sample and is nondiffusable. Particular methods of binding include the use of antibodies (which do not sterically block either the ligand binding site or activation sequence when the protein is bound to the support), direct binding to "sticky" or ionic supports, chemical crosslinking, the synthesis of the protein or agent on the surface, etc. Following binding of the sample, excess unbound material is removed by washing. The sample receiving areas may then be blocked through incubation with bovine

serum albumin (BSA), casein or other innocuous protein or other moiety.

[00155] The compounds of the invention may be used on their own to inhibit the activity of a mitotic kinesin, particularly KSP. In one embodiment, a compound of the invention is combined with KSP and the activity of KSP is assayed. Kinesin (including KSP) activity is known in the art and includes one or more kinesin activities. Kinesin activities include the ability to affect ATP hydrolysis; microtubule binding; gliding and polymerization/depolymerization (effects on microtubule dynamics); binding to other proteins of the spindle; binding to proteins involved in cell-cycle control; serving as a substrate to other enzymes, such as kinases or proteases; and specific kinesin cellular activities such as spindle pole separation.

[00156] Methods of performing motility assays are well known to those of skill in the art. (See e.g., Hall, et al. (1996), *Biophys. J.*, 71: 3467-3476, Turner et al., 1996, *Anal Biochem.* 242 (1):20-5; Gittes et al., 1996, *Biophys. J.* 70(1): 418-29; Shirakawa et al., 1995, *J. Exp. Biol.* 198: 1809-15; Winkelmann et al., 1995, *Biophys. J.* 68: 2444-53; Winkelmann et al., 1995, *Biophys. J.* 68: 72S.)

[00157] Methods known in the art for determining ATPase hydrolysis activity also can be used. Suitably, solution based assays are utilized. U.S. Patent 6,410,254, hereby incorporated by reference in its entirety, describes such assays. Alternatively, conventional methods are used. For example, P_i release from kinesin can be quantified. In one embodiment, the ATPase hydrolysis activity assay utilizes 0.3 M PCA (perchloric acid) and malachite green reagent (8.27 mM sodium molybdate II, 0.33 mM malachite green oxalate, and 0.8 mM Triton X-100). To perform the assay, 10 μ L of the reaction mixture is quenched in 90 μ L of cold 0.3 M PCA. Phosphate standards are used so data can be converted to mM inorganic phosphate released. When all reactions and standards have been quenched in PCA, 100 μ L of malachite green reagent is added to the relevant wells in e.g., a microtiter plate. The mixture is developed for 10-15 minutes and the plate is read at an absorbance of 650 nm. If phosphate standards were used, absorbance readings can be converted to mM P_i and plotted over time. Additionally, ATPase assays known in the art include the luciferase assay.

[00158] ATPase activity of kinesin motor domains also can be used to monitor the effects of agents and are well known to those skilled in the art. In one embodiment ATPase assays of kinesin are performed in the absence of microtubules.

In another embodiment, the ATPase assays are performed in the presence of microtubules. Different types of agents can be detected in the above assays. In a one embodiment, the effect of an agent is independent of the concentration of microtubules and ATP. In another embodiment, the effect of the agents on kinesin ATPase can be decreased by increasing the concentrations of ATP, microtubules or both. In yet another embodiment, the effect of the agent is increased by increasing concentrations of ATP, microtubules or both.

[00159] Compounds that inhibit the biochemical activity of KSP in vitro may then be screened in vivo. In vivo screening methods include assays of cell cycle distribution, cell viability, or the presence, morphology, activity, distribution, or number of mitotic spindles. Methods for monitoring cell cycle distribution of a cell population, for example, by flow cytometry, are well known to those skilled in the art, as are methods for determining cell viability. See for example, U.S. Patent 6,437,115, hereby incorporated by reference in its entirety. Microscopic methods for monitoring spindle formation and malformation are well known to those of skill in the art (see, e.g., Whitehead and Rattner (1998), J. Cell Sci. 111:2551-61; Galgio et al, (1996) J. Cell Biol., 135:399-414), each incorporated herein by reference in its entirety.

[00160] The compounds of the invention inhibit the KSP kinesin. One measure of inhibition is IC_{50} , defined as the concentration of the compound at which the activity of KSP is decreased by fifty percent relative to a control. Preferred compounds have IC_{50} 's of less than about 1 mM, with preferred embodiments having IC_{50} 's of less than about 100 μ M, with more preferred embodiments having IC_{50} 's of less than about 10 μ M, with particularly preferred embodiments having IC_{50} 's of less than about 1 μ M, and especially preferred embodiments having IC_{50} 's of less than about 100 nM, and with the most preferred embodiments having IC_{50} 's of less than about 10 nM. Measurement of IC_{50} is done using an ATPase assay such as described herein.

[00161] Another measure of inhibition is K_i . For compounds with IC_{50} 's less than 1 μ M, the K_i or K_d is defined as the dissociation rate constant for the interaction of the compounds described herein with KSP. Preferred compounds have K_i 's of less than about 100 μ M, with preferred embodiments having K_i 's of less than about 10 μ M, and particularly preferred embodiments having K_i 's of less than about 1 μ M and

especially preferred embodiments having K_i 's of less than about 100 nM, and with the most preferred embodiments having K_i 's of less than about 10 nM.

[00162] The K_i for a compound is determined from the IC_{50} based on three assumptions and the Michaelis-Menten equation. First, only one compound molecule binds to the enzyme and there is no cooperativity. Second, the concentrations of active enzyme and the compound tested are known (i.e., there are no significant amounts of impurities or inactive forms in the preparations). Third, the enzymatic rate of the enzyme-inhibitor complex is zero. The rate (i.e., compound concentration) data are fitted to the equation:

$$V = V_{\max} E_0 \left[I - \frac{(E_0 + I_0 + K_d) - \sqrt{(E_0 + I_0 + K_d)^2 - 4 E_0 I_0}}{2 E_0} \right]$$

where V is the observed rate, V_{\max} is the rate of the free enzyme, I_0 is the inhibitor concentration, E_0 is the enzyme concentration, and K_d is the dissociation constant of the enzyme-inhibitor complex.

[00163] Another measure of inhibition is GI_{50} , defined as the concentration of the compound that results in a decrease in the rate of cell growth by fifty percent. Preferred compounds have GI_{50} 's of less than about 1 mM; those having a GI_{50} of less than about 20 μ M are more preferred; those having a GI_{50} of less than about 10 μ M more so; those having a GI_{50} of less than about 1 μ M more so; those having a GI_{50} of less than about 100 nM more so; and those having a GI_{50} of less than about 10 nM even more so. Measurement of GI_{50} is done using a cell proliferation assay such as described herein. Compounds of this class were found to inhibit cell proliferation.

[00164] In vitro potency of small molecule inhibitors is determined, for example, by assaying human ovarian cancer cells (SKOV3) for viability following a 72-hour exposure to a 9-point dilution series of compound. Cell viability is determined by measuring the absorbance of formazon, a product formed by the bio-reduction of MTS/PMS, a commercially available reagent. Each point on the dose-response curve is calculated as a percent of untreated control cells at 72 hours minus background absorption (complete cell kill).

[00165] Anti-proliferative compounds that have been successfully applied in the clinic to treatment of cancer (cancer chemotherapeutics) have GI_{50} 's that vary

greatly. For example, in A549 cells, paclitaxel GI_{50} is 4 nM, doxorubicin is 63 nM, 5-fluorouracil is 1 μ M, and hydroxyurea is 500 μ M (data provided by National Cancer Institute, Developmental Therapeutic Program, <http://dtp.nci.nih.gov/>). Therefore, compounds that inhibit cellular proliferation, irrespective of the concentration demonstrating inhibition, have potential clinical usefulness.

[00166] To employ the compounds of the invention in a method of screening for compounds that bind to KSP kinesin, the KSP is bound to a support, and a compound of the invention is added to the assay. Alternatively, the compound of the invention is bound to the support and KSP is added. Classes of compounds among which novel binding agents may be sought include specific antibodies, non-natural binding agents identified in screens of chemical libraries, peptide analogs, etc. Of particular interest are screening assays for candidate agents that have a low toxicity for human cells. A wide variety of assays may be used for this purpose, including labeled in vitro protein-protein binding assays, electrophoretic mobility shift assays, immunoassays for protein binding, functional assays (phosphorylation assays, etc.) and the like.

[00167] The determination of the binding of the compound of the invention to KSP may be done in a number of ways. In one embodiment, the compound is labeled, for example, with a fluorescent or radioactive moiety, and binding is determined directly. For example, this may be done by attaching all or a portion of KSP to a solid support, adding a labeled test compound (for example a compound of the invention in which at least one atom has been replaced by a detectable isotope), washing off excess reagent, and determining whether the amount of the label is that present on the solid support.

[00168] By "labeled" herein is meant that the compound is either directly or indirectly labeled with a label which provides a detectable signal, e.g., radioisotope, fluorescent tag, enzyme, antibodies, particles such as magnetic particles, chemiluminescent tag, or specific binding molecules, etc. Specific binding molecules include pairs, such as biotin and streptavidin, digoxin and antidigoxin etc. For the specific binding members, the complementary member would normally be labeled with a molecule which provides for detection, in accordance with known procedures, as outlined above. The label can directly or indirectly provide a detectable signal.

[00169] In some embodiments, only one of the components is labeled. For

example, the kinesin proteins may be labeled at tyrosine positions using ^{125}I , or with fluorophores. Alternatively, more than one component may be labeled with different labels; using ^{125}I for the proteins, for example, and a fluorophor for the antimitotic agents.

[00170] The compounds of the invention may also be used as competitors to screen for additional drug candidates. "Candidate agent" or "drug candidate" or grammatical equivalents as used herein describe any molecule, e.g., protein, oligopeptide, small organic molecule, polysaccharide, polynucleotide, etc., to be tested for bioactivity. They may be capable of directly or indirectly altering the cellular proliferation phenotype or the expression of a cellular proliferation sequence, including both nucleic acid sequences and protein sequences. In other cases, alteration of cellular proliferation protein binding and/or activity is screened. Screens of this sort may be performed either in the presence or absence of microtubules. In the case where protein binding or activity is screened, particular embodiments exclude molecules already known to bind to that particular protein, for example, polymer structures such as microtubules, and energy sources such as ATP. Particular embodiments of assays herein include candidate agents which do not bind the cellular proliferation protein in its endogenous native state termed herein as "exogenous" agents. In another embodiment, exogenous agents further exclude antibodies to KSP.

[00171] Candidate agents can encompass numerous chemical classes, though typically they are organic molecules, preferably they are small organic compounds having a molecular weight of more than 100 and less than about 2,500 daltons. Candidate agents comprise functional groups necessary for structural interaction with proteins, particularly hydrogen bonding and lipophilic binding, and typically include at least an amine, carbonyl-, hydroxyl-, ether, or carboxyl group, preferably at least two of the functional chemical groups. The candidate agents often comprise cyclical carbon or heterocyclic structures and/or aromatic or polyaromatic structures substituted with one or more of the above functional groups. Candidate agents are also found among biomolecules including peptides, saccharides, fatty acids, steroids, purines, pyrimidines, derivatives, structural analogs or combinations thereof.

[00172] Candidate agents are obtained from a wide variety of sources including libraries of synthetic or natural compounds. For example, numerous means are available for random and directed synthesis of a wide variety of organic compounds

and biomolecules, including expression of randomized oligonucleotides. Alternatively, libraries of natural compounds in the form of bacterial, fungal, plant and animal extracts are available or readily produced. Additionally, natural or synthetically produced libraries and compounds are readily modified through conventional chemical, physical and biochemical means. Known pharmacological agents may be subjected to directed or random chemical modifications, such as acylation, alkylation, esterification, and/or amidification to produce structural analogs.

[00173] Competitive screening assays may be done by combining KSP and a drug candidate in a first sample. A second sample comprises a compound of the present invention, KSP and a drug candidate. This may be performed in either the presence or absence of microtubules. The binding of the drug candidate is determined for both samples, and a change, or difference in binding between the two samples indicates the presence of a drug candidate capable of binding to KSP and potentially inhibiting its activity. That is, if the binding of the drug candidate is different in the second sample relative to the first sample, the drug candidate is capable of binding to KSP.

[00174] In a particular embodiment, the binding of the candidate agent to KSP is determined through the use of competitive binding assays. In this embodiment, the competitor is a binding moiety known to bind to KSP, such as an antibody, peptide, binding partner, ligand, etc. Under certain circumstances, there may be competitive binding as between the candidate agent and the binding moiety, with the binding moiety displacing the candidate agent.

[00175] In one embodiment, the candidate agent is labeled. Either the candidate agent, or the competitor, or both, is added first to KSP for a time sufficient to allow binding, if present. Incubations may be performed at any temperature which facilitates optimal activity, typically between 4 and 40°C.

[00176] Incubation periods are selected for optimum activity, but may also be optimized to facilitate rapid high throughput screening. Typically between 0.1 and 1 hour will be sufficient. Excess reagent is generally removed or washed away. The second component is then added, and the presence or absence of the labeled component is followed, to indicate binding.

[00177] In another embodiment, the competitor is added first, followed by the candidate agent. Displacement of the competitor is an indication the candidate agent

is binding to KSP and thus is capable of binding to, and potentially inhibiting, the activity of KSP. In this embodiment, either component can be labeled. Thus, for example, if the competitor is labeled, the presence of label in the wash solution indicates displacement by the agent. Alternatively, if the candidate agent is labeled, the presence of the label on the support indicates displacement.

[00178] In an alternative embodiment, the candidate agent is added first, with incubation and washing, followed by the competitor. The absence of binding by the competitor may indicate the candidate agent is bound to KSP with a higher affinity. Thus, if the candidate agent is labeled, the presence of the label on the support, coupled with a lack of competitor binding, may indicate the candidate agent is capable of binding to KSP.

[00179] Inhibition is tested by screening for candidate agents capable of inhibiting the activity of KSP comprising the steps of combining a candidate agent with KSP, as above, and determining an alteration in the biological activity of KSP. Thus, in this embodiment, the candidate agent should both bind to KSP (although this may not be necessary), and alter its biological or biochemical activity as defined herein. The methods include both in vitro screening methods and in vivo screening of cells for alterations in cell cycle distribution, cell viability, or for the presence, morphology, activity, distribution, or amount of mitotic spindles, as are generally outlined above.

[00180] Alternatively, differential screening may be used to identify drug candidates that bind to the native KSP, but cannot bind to modified KSP.

[00181] Positive controls and negative controls may be used in the assays. Suitably all control and test samples are performed in at least triplicate to obtain statistically significant results. Incubation of all samples is for a time sufficient for the binding of the agent to the protein. Following incubation, all samples are washed free of non-specifically bound material and the amount of bound, generally labeled agent determined. For example, where a radiolabel is employed, the samples may be counted in a scintillation counter to determine the amount of bound compound.

[00182] A variety of other reagents may be included in the screening assays. These include reagents like salts, neutral proteins, e.g., albumin, detergents, etc which may be used to facilitate optimal protein-protein binding and/or reduce non-specific or background interactions. Also reagents that otherwise improve the efficiency of the

assay, such as protease inhibitors, nuclease inhibitors, anti-microbial agents, etc., may be used. The mixture of components may be added in any order that provides for the requisite binding.

Administration

[00183] Accordingly, the compounds of the invention are administered to cells. By "administered" herein is meant administration of a therapeutically effective dose of a compound of the invention to a cell either in cell culture or in a patient. By "therapeutically effective dose" herein is meant a dose that produces the effects for which it is administered. The exact dose will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques. As is known in the art, adjustments for systemic versus localized delivery, age, body weight, general health, sex, diet, time of administration, drug interaction and the severity of the condition may be necessary, and will be ascertainable with routine experimentation by those skilled in the art. By "cells" herein is meant any cell in which mitosis or meiosis can be altered.

[00184] A "patient" for the purposes of the present invention includes both humans and other animals, particularly mammals, and other organisms. Thus the methods are applicable to both human therapy and veterinary applications. In a particular embodiment the patient is a mammal, and more particularly, the patient is human.

[00185] Compounds of the invention having the desired pharmacological activity may be administered, generally as a pharmaceutically acceptable composition comprising an pharmaceutical excipient, to a patient, as described herein. Depending upon the manner of introduction, the compounds may be formulated in a variety of ways as discussed below. The concentration of therapeutically active compound in the formulation may vary from about 0.1-100 wt.%.

[00186] The agents may be administered alone or in combination with other treatments, i.e., radiation, or other chemotherapeutic agents such as the taxane class of agents that appear to act on microtubule formation or the camptothecin class of topoisomerase I inhibitors. When used, other chemotherapeutic agents may be administered before, concurrently, or after administration of a compound of the present invention. In one aspect of the invention, a compound of the present invention

is co-administered with one or more other chemotherapeutic agents. By "co-administer" it is meant that the present compounds are administered to a patient such that the present compounds as well as the co-administered compound may be found in the patient's bloodstream at the same time, regardless when the compounds are actually administered, including simultaneously.

[00187] The administration of the compounds and compositions of the present invention can be done in a variety of ways, including, but not limited to, orally, subcutaneously, intravenously, intranasally, transdermally, intraperitoneally, intramuscularly, intrapulmonary, vaginally, rectally, or intraocularly. In some instances, for example, in the treatment of wounds and inflammation, the compound or composition may be directly applied as a solution or spray.

[00188] Pharmaceutical dosage forms include a compound of formula I or a pharmaceutically acceptable salt, solvate, or solvate of a salt thereof, and one or more pharmaceutical excipients. As is known in the art, pharmaceutical excipients are secondary ingredients which function to enable or enhance the delivery of a drug or medicine in a variety of dosage forms (e.g.: oral forms such as tablets, capsules, and liquids; topical forms such as dermal, ophthalmic, and otic forms; suppositories; injectables; respiratory forms and the like). Pharmaceutical excipients include inert or inactive ingredients, synergists or chemicals that substantively contribute to the medicinal effects of the active ingredient. For example, pharmaceutical excipients may function to improve flow characteristics, product uniformity, stability, taste, or appearance, to ease handling and administration of dose, for convenience of use, or to control bioavailability. While pharmaceutical excipients are commonly described as being inert or inactive, it is appreciated in the art that there is a relationship between the properties of the pharmaceutical excipients and the dosage forms containing them.

[00189] Pharmaceutical excipients suitable for use as carriers or diluents are well known in the art, and may be used in a variety of formulations. See, e.g., Remington's Pharmaceutical Sciences, 18th Edition, A. R. Gennaro, Editor, Mack Publishing Company (1990); Remington: The Science and Practice of Pharmacy, 20th Edition, A. R. Gennaro, Editor, Lippincott Williams & Wilkins (2000); Handbook of Pharmaceutical Excipients, 3rd Edition, A. H. Kibbe, Editor, American Pharmaceutical Association, and Pharmaceutical Press (2000); and Handbook of Pharmaceutical Additives, compiled by Michael and Irene Ash, Gower (1995), each of

which is incorporated herein by reference for all purposes.

[00190] Oral solid dosage forms such as tablets will typically comprise one or more pharmaceutical excipients, which may for example help impart satisfactory processing and compression characteristics, or provide additional desirable physical characteristics to the tablet. Such pharmaceutical excipients may be selected from diluents, binders, glidants, lubricants, disintegrants, colors, flavors, sweetening agents, polymers, waxes or other solubility-retarding materials.

[00191] Compositions for intravenous administration will generally comprise intravenous fluids, i.e., sterile solutions of simple chemicals such as sugars, amino acids or electrolytes, which can be easily carried by the circulatory system and assimilated. Such fluids are prepared with water for injection USP.

[00192] Fluids used commonly for intravenous (IV) use are disclosed in Remington, the Science and Practice of Pharmacy [full citation previously provided], and include:

- alcohol (e.g., in dextrose and water ("D/W") [e.g., 5% dextrose] or dextrose and water [e.g., 5% dextrose] in normal saline solution ("NSS"); e.g. 5% alcohol);
- synthetic amino acid such as Aminosyn, FreAmine, Travasol, e.g., 3.5 or 7; 8.5; 3.5, 5.5 or 8.5 % respectively;
- ammonium chloride e.g., 2.14%;
- dextran 40, in NSS e.g., 10% or in D5/W e.g., 10%;
- dextran 70, in NSS e.g., 6% or in D5/W e.g., 6%;
- dextrose (glucose, D5/W) e.g., 2.5-50%;
- dextrose and sodium chloride e.g., 5-20% dextrose and 0.22-0.9% NaCl;
- lactated Ringer's (Hartmann's) e.g., NaCl 0.6%, KCl 0.03%, CaCl_2 0.02%;
- lactate 0.3%;
- mannitol e.g., 5%, optionally in combination with dextrose e.g., 10% or NaCl e.g., 15 or 20%;
- multiple electrolyte solutions with varying combinations of electrolytes, dextrose, fructose, invert sugar Ringer's e.g., NaCl 0.86%, KCl 0.03%, CaCl_2 0.033%;
- sodium bicarbonate e.g., 5%;
- sodium chloride e.g., 0.45, 0.9, 3, or 5%;
- sodium lactate e.g., 1/6 M; and

sterile water for injection


The pH of such fluids may vary, and will typically be from 3.5 to 8 such as known in the art.


[00193] The following examples serve to more fully describe the manner of using the above-described invention, as well as to set forth the best modes contemplated for carrying out various aspects of the invention. It is understood that these examples in no way serve to limit the true scope of this invention, but rather are presented for illustrative purposes. All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

Examples

Example 1

Synthesis of Compounds





[00194] A solution of 4-chloro-2-iodobenzoic acid 1 (25 g, 88.7 mmol), SOCl_2 (100 mL), and DMF (few drops) was warmed gently with a heat gun until the mixture became homogeneous (15 mins). The solution was maintained at 23°C for an additional 30 mins, then the solution was concentrated. MeOH (200 mL) was added to the crude residue and the solution was maintained at 23°C for 30 mins. The solution was concentrated and the crude residue was then dissolved in 10:1 hexanes:EtOAc and passed through a plug of silica gel. The eluent was concentrated to provide 26.2 g (100%) of ester 2 as a colorless oil, which solidified upon standing under high vacuum (0.1 Torr).

[00195] A mixture of ester 2 (2.35 g, 7.94 mmol), acetylene 31 (1.72 g, 8.7 mmol), $\text{Pd(PPh}_3)_2\text{Cl}_2$ (140 mg, 0.20 mmol), CuI (19 mg, 0.1 mmol), and TEA (35 mL) was maintained at 50°C for 1h. The reaction mixture was diluted with EtOAc (200 mL), washed with water (100 mL), then brine (100 mL). The organic layer was dried (MgSO_4) and concentrated. The crude residue was purified by flash column chromatography (10:1 hexanes:EtOAc) to provide 2.88 g (100%) of 4.

[00196] Ester 4 (2.88 g, 7.94 mmol), KOH (930 mg, 16.5 mmol), MeOH (15 mL), THF (15 mL), and H₂O (15 mL) was maintained at 50°C for exactly 30 mins. The reaction was diluted with EtOAc (100 mL) and washed with 1N HCl (20 mL) and brine (20 mL). The organic layer was dried (MgSO₄) and concentrated to provide 2.76 g of acid 5 (99%), which was used without further purification.

[00197] Pd(MeCN)₂Cl₂ (98mg, 0.39 mmol) was added to a solution of the crude acid 5 (2.76 g, 7.87 mmol), TEA (3.3 mL) and THF (54 mL). The solution was maintained at 50°C for 30 mins. The reaction solution was diluted with hexanes (150 mL) and passed through a plug of silica gel. The plug was rinsed with (5:1 hexanes:EtOAc). The filtrate was concentrated. Hexanes (50 mL) was added to this crude residue and pure 6 precipitated out of solution. This precipitated material was filtered and used without further purification. The filtrate was then concentrated and the remaining residue dried onto silica gel (CH₂Cl₂) and purified by flash column chromatography (10:1 hexanes:EtOAc; 5:1 hexanes:EtOAc; 3:1 hexanes:EtOAc) to provide a combined 2.22 g of 6 (77%, two steps).

[00198] A solution of isochromenone 6 (2.1 g, 6.0 mmol), BnNH₂ (1.96 mL, 17.9 mmol), and PhMe (3.0 mL) was maintained at 140°C for 1.5h. The reaction mixture was cooled to 23°C, diluted with EtOAc (50 mL), washed with 1 N HCl (20 mL) and brine (20 mL). The organic layer was dried (MgSO₄) and concentrated. The resulting crude residue was used without further purification. A solution of the crude residue, 5% HCl (3.4 mL) and MeOH (17 mL) was maintained at 50°C for 1.5 h. The product 7 begins to precipitate from the reaction solution after 5 mins as a white solid. After 30 mins, an additional amount of MeOH (5 mL) and 5% HCl (1 mL) was added and the reaction mixture was maintained for an additional 1h. The reaction mixture was diluted with EtOAc (100 mL), washed with 1 N NaOH (50 mL) and then brine (20 mL). The organic layer was dried (MgSO₄) and concentrated. The residue was purified by flash column chromatography (5:1 hexanes:EtOAc) to provide 1.8 g of racemic 7 (68%, two steps).

[00199] A solution of isoquinolone 7 (2.50 g, 5.68 mmol) and TFA:H₂O (97.5:2.5; 25 mL) was maintained at 23°C for 1h. The solution was concentrated. The crude residue was diluted with EtOAc (75 mL), washed with 1 N NaOH (20 mL), and

1 Fehrentz, J.-A.; Castro, B. *Synthesis* 1983, 676. Goel, O.; Krolls, U.; Steir, M.; Kesten, S. *Org. Syn.*


brine (20 mL). The organic layer was dried (MgSO_4) and concentrated to give 1.83 g of amine **9** (95%) as a white solid.


[00200] Isoquinolone **8** (131 mg, 0.34 mmol), aldehyde **9** (115 mg, 0.56 mmol), $\text{Na}(\text{OAc})_3\text{BH}$ (291 mg, 1.38 mmol), and CH_2Cl_2 (1.1 mL) was maintained at 23°C for 3 h. The reaction mixture was diluted with EtOAc (20 mL) and washed with 1 N NaOH (5 mL) and brine (5 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue was purified by flash column chromatography (3:1 hexanes:EtOAc) to yield 150 mg (85%) of **10**.

[00201] A solution of isoquinolone **10** (143 mg, 0.28 mmol), POCl_3 (0.45 mL, 4.8 mmol), and PhMe (14 mL) was heated to 110°C . After 6h, the reaction mixture was diluted with EtOAc (40 mL) and washed with 1 N NaOH (20 mL) and brine (10 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue was purified by flash column chromatography (20:1 CHCl_3 :MeOH) to yield 99 mg (70%) of **11** as a white solid.

Example 2

Synthesis of Compounds





[00202] Isoquinolone 8 (515 mg, 1.47 mmol), aldehyde 12 (255 mg, 1.47 mmol), $\text{NaCN}(\text{OAc})_3\text{BH}$ (420 mg, 1.98 mmol), and CH_2Cl_2 (4.1 mL) was maintained at 23°C for 2 h. An additional portion of 12 (225 mg, 1.30 mmol) in CH_2Cl_2 (0.6 mL) was then added. After an additional 3h, the reaction mixture was diluted with EtOAc (20 mL) and washed with 1 N NaOH (5 mL) and brine (5 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue was purified by flash column chromatography (3:1 hexanes:EtOAc; 1:1 hexanes:EtOAc) to yield 630 mg (86%) of 13.




[00203] To a solution of isoquinolone 13 (85 mg, 0.17 mmol), diisopropylethylamine (DIEA, 0.12 mL, 0.68 mmol), and CH_2Cl_2 (0.6 mL) at 23°C was added p -toluoyl chloride (45 μL , 0.34 mmol). After 4 h, the reaction mixture was diluted with EtOAc (20 mL) and washed with saturated aqueous NaHCO_3 (5 mL) and brine (5 mL). The organic layer was dried (MgSO_4), filtered, and concentrated. The resulting residue was purified by flash column chromatography (3:1 hexanes:EtOAc) to yield 83 mg (80%) of 14.





[00204] Isoquinolone 14 (80 mg, 0.13 mmol) and TFA: H_2O (97.5:2.5, 2 mL) was maintained at 23°C for 1h. The reaction mixture was concentrated. The residue was dissolved in EtOAc (20 mL) and washed with 1 N NaOH (5 mL) and brine (5





mL). The organic layer was dried (MgSO_4), filtered, and concentrated to provide 65 mg (98%) of **15** as a white solid which was deemed >95% pure by ^1H NMR and LCMS analysis.


Example 3

[00205] Using the methods of the invention as exemplified in Examples 1 and 2 above, the following compounds were prepared:

Structure	LRMS (MH) m/z
	516.2
	470.2
	512.2

	575.8
	528.0
	464.0
	530.0

	516.2
	566.2
	512.2
	565.8



Example 4

Monopolar Spindle Formation following Application of an Isoquinolone KSP

Inhibitor

[00206] Human tumor cell lines Skov-3 (ovarian) are plated in 96-well plates at densities of 4,000 cells per well, allowed to adhere for 24 hours, and treated with various concentrations of the isoquinolone compounds for 24 hours. Cells are fixed in 4% formaldehyde and stained with antitubulin antibodies (subsequently recognized using fluorescently-labeled secondary antibody) and Hoechst dye (which stains DNA).

[00207] Visual inspection reveals that the compounds cause cell cycle arrest in the prometaphase stage of mitosis. DNA is condensed and spindle formation is initiated, but arrested cells uniformly display monopolar spindles, indicating that there is an inhibition of spindle pole body separation. Microinjection of anti-KSP antibodies also causes mitotic arrest with arrested cells displaying monopolar spindles.

Example 5

Inhibition of Cellular Proliferation in Tumor Cell Lines Treated with KSP

Inhibitors.

[00208] Cells are plated in 96-well plates at densities from 1000-2500 cells/well of a 96-well plate and allowed to adhere/grow for 24 hours. They are then treated with various concentrations of drug for 48 hours. The time at which compounds are added is considered T_0 . A tetrazolium-based assay using the reagent 3-(4,5-dimethylthiazol-2-yl)-5-(3-carboxymethoxyphenyl)-2-(4-sulfophenyl)-2H-tetrazolium (MTS) (I.S. Patent No. 5,185,450) (see Promega product catalog #G3580,

CellTiter 96® AQueous One Solution Cell Proliferation Assay) is used to determine the number of viable cells at T_0 and the number of cells remaining after 48 hours compound exposure. The number of cells remaining after 48 hours is compared to the number of viable cells at the time of drug addition, allowing for calculation of growth inhibition.

[00209] The growth over 48 hours of cells in control wells that have been treated with vehicle only (0.25% DMSO) is considered 100% growth and the growth of cells in wells with compounds is compared to this. KSP inhibitors inhibit cell proliferation in human ovarian tumor cell lines (SKOV-3).

[00210] A GI_{50} is calculated by plotting the concentration of compound in μM vs the percentage of cell growth of cell growth in treated wells. The GI_{50} calculated for the compounds is the estimated concentration at which growth is inhibited by 50% compared to control, i.e., the concentration at which:

$$100 \times [(Treated_{48} - T_0) / (Control_{48} - T_0)] = 50.$$

[00211] All concentrations of compounds are tested in duplicate and controls are averaged over 12 wells. A very similar 96-well plate layout and GI_{50} calculation scheme is used by the National Cancer Institute (see Monks, et al., J. Natl. Cancer Inst. 83:757-766 (1991)). However, the method by which the National Cancer Institute quantitates cell number does not use MTS, but instead employs alternative methods.

Example 6

Calculation of IC_{50} :

[00212] Measurement of a compound's IC_{50} for KSP activity uses an ATPase assay. The following solutions are used: Solution 1 consists of 3 mM phosphoenolpyruvate potassium salt (Sigma P-7127), 2 mM ATP (Sigma A-3377), 1 mM IDTT (Sigma D-9779), 5 μM paclitaxel (Sigma T-7402), 10 ppm antifoam 289 (Sigma A-8436), 25 mM Pipes/KOH pH 6.8 (Sigma P6757), 2 mM $MgCl_2$ (VWR JT400301), and 1 mM EGTA (Sigma E3889). Solution 2 consists of 1 mM NADH (Sigma N8129), 0.2 mg/ml BSA (Sigma A7906), pyruvate kinase 7U/ml, L-lactate dehydrogenase 10 U/ml (Sigma P0294), 100 nM KSP motor domain, 50 $\mu g/ml$ microtubules, 1 mM DTT (Sigma D9779), 5 μM paclitaxel (Sigma T-7402), 10 ppm antifoam 289 (Sigma A-8436), 25 mM Pipes/KOH pH 6.8 (Sigma P6757), 2 mM


MgC12 (VWR JT4003-01), and 1 mM EGTA (Sigma E3889). Serial dilutions (8-12 two-fold dilutions) of the compounds are made in a 96-well microtiter plate (Corning Costar 3695) using Solution 1. Following serial dilution each well has 50 μ l of Solution 1. The reaction is started by adding 50 μ l of solution 2 to each well. This may be done with a multichannel pipettor either manually or with automated liquid handling devices. The microtiter plate is then transferred to a microplate absorbance reader and multiple absorbance readings at 340 nm are taken for each well in a kinetic mode. The observed rate of change, which is proportional to the ATPase rate, is then plotted as a function of the compound concentration. For a standard IC₅₀ determination the data acquired is fit by the following four parameter equation using a nonlinear fitting program (e.g., Grafit 4):

$$y = \frac{\text{Range}}{1 + \left(\frac{x}{\text{IC}_{50}} \right)^s} + \text{Background}$$

where y is the observed rate and x the compound concentration.

What is claimed is:

1. A compound having the structure represented by Formula I:



(I)

wherein:

R_1 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-;

R_2 and $R_{2'}$ are independently chosen from hydrogen, optionally substituted alkyl-, optionally substituted alkoxy, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-; or R_2 and $R_{2'}$ taken together form an optionally substituted 3- to 7-membered ring;

R_{12} is selected from the group consisting of optionally substituted imidazolyl-, optionally substituted imidazolyl-, $-NHR_4$; $-N(R_4)(COR_3)$; $-N(R_4)(SO_2R_{3a})$; and $-N(R_4)(CH_2R_{3b})$;

R_3 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, optionally substituted heteroaralkyl-, $R_{15}O-$ and $R_{17}NH-$;

R_{3a} is chosen from optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, optionally substituted heteroaralkyl-, and $R_{17}NH-$;

R_{3b} is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-;

R₄ is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heterocycl-, and optionally substituted heteroaralkyl-;

R₅, R₆, R₇ and R₈ are independently chosen from hydrogen, optionally substituted alkyl, optionally substituted alkoxy, halogen, hydroxyl, nitro, cyano, dialkylamino, alkylsulfonyl, alkylsulfonamido, alkylthio, carboxyalkyl, carboxamido, aminocarbonyl, optionally substituted aryl and optionally substituted heteroaryl;

R₁₅ is optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, or optionally substituted heteroaralkyl-; and

R₁₇ is chosen from hydrogen, optionally substituted alkyl, optionally substituted aryl, optionally substituted aralkyl, optionally substituted heteroaryl, and optionally substituted heteroaralkyl;

a pharmaceutically acceptable salt of a compound of Formula I;

a pharmaceutically acceptable solvate of a compound of Formula I;

or a pharmaceutically acceptable solvate of a pharmaceutically acceptable salt of a compound of Formula I.

2. A compound according to claim 1, wherein R₁ is selected from hydrogen, optionally substituted C₁-C₈ alkyl-, optionally substituted aryl-, optionally substituted heteroaryl-, optionally substituted aryl-C₁-C₄-alkyl-, and optionally substituted heteroaryl-C₁-C₄-alkyl-.

3. A compound according to claim 2, wherein R₁ is selected from hydrogen, optionally substituted C₁-C₄ alkyl-, optionally substituted phenyl-C₁-C₄-alkyl-, optionally substituted naphthalenylmethyl-, optionally substituted phenyl-, and naphthyl-.

4. A compound according to claim 3, wherein R₁ is optionally substituted phenyl-C₁-C₄-alkyl-, optionally substituted heteroaryl-C₁-C₄-alkyl-, or

naphthalenylmethyl-.

5. A compound according to claim 4, wherein R_1 is naphthyl-, phenyl-, bromophenyl-, chlorophenyl-, methoxyphenyl-, ethoxyphenyl-, tolyl-, dimethylphenyl-, chlorofluorophenyl-, methylchlorophenyl-, ethylphenyl-, phenethyl-, benzyl-, chlorobenzyl-, methylbenzyl-, methoxybenzyl-, cyanobenzyl-, hydroxybenzyl-, dichlorobenzyl-, dimethoxybenzyl-, or naphthalenylmethyl-.

6. A compound according to claim 5, wherein R_1 is benzyl.

7. A compound according to claim 1, wherein R_2 and R_2' are independently chosen from hydrogen, optionally substituted alkyl-, optionally substituted alkoxy, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaryl-, and optionally substituted heteroaralkyl-; or R_2 and R_2' taken together form an optionally substituted 3- to 7-membered ring.

8. A compound according to claim 7, wherein R_2 is optionally substituted C_1 - C_4 alkyl-, and R_2' is hydrogen or optionally substituted C_1 - C_4 alkyl-.

9. A compound according to claim 8, wherein R_2 is hydrogen and R_2' is optionally substituted C_1 - C_4 alkyl-.

10. A compound according to claim 9, wherein R_2 is hydrogen and R_2' is ethyl or propyl.

11. A compound according to claim 10, wherein R_2 is i-propyl.


12. A compound according to claim 1, wherein if either R_2 or R_2' is hydrogen, then the other is not hydrogen.

13. A compound according to claim 1, wherein R_5 , R_6 , R_7 , and R_8 are independently chosen from hydrogen, hydroxyl, halogen, optionally substituted C_1 - C_4 alkyl-, C_1 - C_4 alkoxy, cyano, amino, substituted amino, or carbamyl-.

14. A compound according to claim 13, wherein R_5 , R_6 , R_7 , and R_8 are independently methoxy, methyl, trifluoromethyl, cyano, hydrogen or halo.

15. A compound according to claim 14, wherein only one of R_5 , R_6 , R_7 , and R_8 is not hydrogen.

16. A compound according to claim 1, wherein R_{12} is an optionally substituted imidazolyl- having the formula:



wherein


R_9 is chosen from hydrogen, optionally substituted C_1 - C_8 alkyl-, optionally substituted aryl-, optionally substituted aryl- C_1 - C_4 -alkyl-, optionally substituted heteroaryl- C_1 - C_4 -alkyl-, optionally substituted aryl- C_1 - C_4 -alkoxy, optionally substituted heteroaryl- C_1 - C_4 -alkoxy, and optionally substituted heteroaryl-; and

R_{13} and $R_{13'}$ are independently hydrogen, optionally substituted C_1 - C_8 alkyl-, optionally substituted aryl-, or optionally substituted aryl- C_1 - C_4 -alkyl-.

17. A compound according to claim 16, wherein R_9 is lower-alkyl; phenyl-; or phenyl- substituted with one or more of the following groups: methyl, methoxy, trifluoromethyl, or halo.

18. A compound according to claim 16, wherein R_{13} is hydrogen and $R_{13'}$ is optionally substituted substituted C_1 - C_4 alkyl-.

19. A compound according to claim 1, wherein R_{12} is an optionally substituted imidazoline having the formula



wherein,

R_{14} is chosen from hydrogen, optionally substituted C_1 - C_8 alkyl-, optionally substituted aryl-, optionally substituted aryl- C_1 - C_4 -alkyl-, optionally substituted heteroaryl-, optionally substituted heteroaryl- C_1 - C_4 -alkyl-; and

R_{10} , $R_{10'}$, R_{11} and $R_{11'}$ are independently chosen from hydrogen, optionally substituted C_1 - C_8 alkyl-, optionally substituted aryl-, and optionally substituted aryl- C_1 - C_4 -alkyl-.

20. A compound according to claim 19, wherein R_{10} , $R_{10'}$, $R_{11'}$, and R_{11} are independently hydrogen or optionally substituted C_1 - C_4 alkyl-.

21. A compound according to claim 1 wherein R_{12} is $-NHR_4$, $-NR_4(COR_3)$, $NR_4(SO_2R_{3a})$, or $-NR_4(CH_2R_{3b})$ and R_4 is R_{16} -alkylene-, and R_{16} is chosen from alkoxy, amino, alkylamino, dialkylamino, carboxy, guanidine, hydroxyl-, and N-heterocycl-.

22. A compound according to claim 1 wherein R_{12} is $-NR_4(COR_3)$ and R_3 is selected from hydrogen, optionally substituted alkyl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, optionally substituted heteroaryl-, optionally substituted aryl-, $R_{15}O$ - and $R_{17}NH$ -, wherein R_{15} is chosen from optionally substituted alkyl and optionally substituted aryl and R_{17} is chosen from hydrogen, optionally substituted alkyl and optionally substituted aryl-.

23. A compound according to claim 22 wherein R_{12} is $-NR_4(COR_3)$ and R_3 is tolyl-, halophenyl-, halomethylphenyl-, hydroxymethylphenyl-, methylenedioxyphenyl-, formylphenyl or cyanophenyl-.

24. A compound according to claim 21 wherein R_{3a} is chosen from C_1 - C_{13} alkyl-; phenyl-; naphthyl-; phenyl substituted with cyano, halo, lower-alkyl-, lower-alkoxy, nitro, methylenedioxy, or trifluoromethyl-; biphenyl and heteroaryl-.

25. A compound according to claim 21 wherein R_{3b} is chosen from C_1 - C_{13} alkyl-; substituted lower-alkyl-; phenyl-; naphthyl-; phenyl substituted with cyano, halo, lower-alkyl-, lower-alkoxy, nitro, methylenedioxy, or trifluoromethyl-; biphenyl-, benzyl and heterocycl-.

26. A compound according to claim 1 wherein,

R_1 is optionally substituted aryl- C_1 - C_4 -alkyl-, optionally substituted heteroaryl- C_1 - C_4 -alkyl-, or naphthalenylmethyl ;

R_2 is optionally substituted C_1 - C_4 -alkyl-;

R_2 is hydrogen;


R_7 is hydrogen, methyl, or cyano;

R_5 , R_6 , and R_8 are hydrogen;

and

R_{12} is optionally substituted imidazolyl-, optionally substituted imidazolyl-, $-NHR_4$; $-N(R_4)(COR_3)$; $-N(R_4)(SO_2 R_{3a})$; and $-N(R_4)(CH_2 R_{3b})$.

27. A compound according to claim 26 wherein R_{12} is optionally substituted imidazolyl- having the formula:




wherein

R_9 is lower-alkyl; phenyl-; or phenyl- substituted with one or more of the following groups: methyl, methoxy, trifluoromethyl, or halo;

R_{13} is hydrogen; and

R_{13} is substituted C_1 - C_4 alkyl-.

28. A compound according to claim 26 wherein R_{12} is optionally substituted imidazoliny- having the formula:



wherein

R_{14} is lower-alkyl; phenyl-; or phenyl- substituted with one or more of the following groups: methyl, methoxy, trifluoromethyl, or halo; and

R_{10} , $R_{10'}$, $R_{11'}$, and R_{11} are independently hydrogen or optionally substituted C_1 - C_4 alkyl-.

29. A compound according to claim 26 wherein R_{12} is $-NHR_4$; and R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, and optionally substituted heterocycl-.

30. A compound according to claim 26 wherein R_3 is selected from hydrogen, optionally substituted alkyl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, optionally substituted heteroaryl-, optionally substituted aryl-, $R_{15}O$ - and $R_{17}NH$ -, wherein R_{15} is chosen from optionally substituted alkyl and optionally substituted aryl and R_{17} is chosen from hydrogen, optionally substituted alkyl and optionally substituted aryl.

31. A compound according to claim 30 wherein R_4 is R_{16} -alkylene-, and R_{16} is chosen from alkoxy, amino, alkylamino, dialkylamino, carboxy, hydroxyl-, and N-heterocycl-.

32. A compound according to claim 31, wherein

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, and optionally substituted heterocyclyl- and

R_3 is selected from optionally substituted alkyl-, aryl-, substituted aryl-, benzyl-, and optionally substituted heteroaryl-.

33. A compound according to claim 32, wherein R_3 is tolyl-, halophenyl-, halomethylphenyl-, hydroxymethylphenyl-, methylenedioxyphenyl-, formylphenyl or cyanophenyl-.

34. A compound according to claim 26, wherein R_{12} is $-N(R_4)(CH_2 R_{3b})$;

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, and optionally substituted heterocyclyl- and

R_{3b} is chosen from phenyl substituted with one or more halo, methyl-, cyano, trifluoromethyl-, trifluoromethoxy, carboxy, or methoxycarbonyl groups; piperidinyl-; and naphthyl-.

35. A compound according to claim 1, wherein R_{12} is $-NR_4(SO_2 R_{3a})$;

R_4 is chosen from hydrogen, optionally substituted alkyl-, optionally substituted aryl-, optionally substituted aralkyl-, optionally substituted heteroaralkyl-, and optionally substituted heterocyclyl- and

R_{3a} is chosen from phenyl substituted with halo, lower-alkyl-, lower-alkoxy, cyano, nitro, methylenedioxy, or trifluoromethyl-; and naphthyl-.

36. A compound according to any one of claims 1-35 wherein R_2 and R_2' are each attached to a stereogenic center having an R-configuration.

37. A composition comprising a pharmaceutical excipient and a compound, salt, or solvate thereof of any one of claims 1-36.

38. A composition according to claim 37, wherein said composition further

comprises a chemotherapeutic agent other than a compound of Formula I or said salt, solvate, or solvate of a salt thereof.

39. A composition according to claim 38, wherein said composition further comprises a taxane.

40. A composition according to claim 38, wherein said composition further comprises a vinca alkaloid.

41. A composition according to claim 38, wherein said composition further comprises a topoisomerase I inhibitor.

42. A method of modulating KSP kinesin activity which comprises contacting said kinesin with an effective amount of a compound according to any one of claims 1 to 41.

43. A method of inhibiting KSP which comprises contacting said kinesin with an effective amount of a compound according to any one of claims 1 to 41.

44. A method for the treatment of a cellular proliferative disease comprising administering to a subject in need thereof a compound according to any one of claims 1-41.

45. A method for the treatment of a cellular proliferative disease comprising administering to a subject in need thereof a composition according to any one of claims 42-44.

46. A method according to claim 44 or claim 45 wherein said disease is selected from the group consisting of cancer, hyperplasias, restenosis, cardiac hypertrophy, immune disorders, and inflammation.

47. The use, in the manufacture of a medicament for treating cellular proliferative disease, of a compound according to any one of claims 1-41.

48. The use of a compound as defined in claim 47 for the manufacture of a medicament for treating a disorder associated with KSP kinesin activity.